

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 April 2005 (07.04.2005)

PCT

(10) International Publication Number
WO 2005/030786 A1

(51) International Patent Classification⁷: C07H 17/08,
A61K 31/7048

(21) International Application Number:
PCT/IB2003/004166

(22) International Filing Date:
25 September 2003 (25.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): RAN-
BAXY LABORATORIES LIMITED [IN/IN]; 19, Nehru
Place, New Delhi, Delhi 110 019 (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DAS, Biswajit
[IN/IN]; 2nd Floor, J7/13 DLF II, Gurgaon, Haryana 122
001 (IN). SALMAN, Mohammad, [IN/IN]; P - 2/19,
DLF City, Phase - II, Gurgaon, Haryana 122 001 (IN).
SHELKE, Sachin, Vilasrao [IN/IN]; Gautam Public
School, Kolpewadi, Tal-kopargaon, Ahmadnagar, Maha-
rastra 423 601 (IN). HAJARE, Atul, Kashinath [IN/IN];
House No. 700, Sector 17A, Gurgaon, Haryana, 122 001
(IN). TALUKDAR, Sanjay [IN/IN]; 64/3, Ashokgarh,
Kolkata, West Bengal 700 035 (IN). RATHY, Sujata

[IN/IN]; WZ-325 Rishi Nagar, Rani Bagh, Delhi, Delhi
110 034 (IN). PAL, Arani [IN/IN]; Basundhara Apartment
6E, 19/922, Raja S.C. Mullick Road, Kolkata, West Bengal
700 086 (IN). RATTAN; Ashok [IN/IN]; B-481 Sarita
Vihar, New Dehli, Delhi 110 044 (IN).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KB, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 3'-N-SUBSTITUTED-3-O-SUBSTITUTED ERYTHRONOLIDE A DERIVATIVES

(57) Abstract: This invention generally relates to macrolides, more particularly, the invention relates to 3-N-substituted-3-O-substituted erythronolide A derivatives, which are antibacterial agents effective against gram positive or gram negative bacteria and atypical pathogens. The compounds of this invention are more particularly effective against *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Haemophilus influenzae*. The invention also relates to a process for the preparation of the compounds of the present invention, pharmaceutical compositions containing the compounds of the present invention and the methods for treating bacterial infection.

3'-N-SUBSTITUTED-3-O-SUBSTITUTED ERYTHRONOLIDE A DERIVATIVES**FIELD OF INVENTION**

This invention generally relates to macrolides, more particularly, the invention relates to 3'-N-substituted-3-O-substituted erythronolide A derivatives, which are antibacterial agents effective against gram positive or gram negative bacteria and atypical pathogens. The compounds of this invention are more particularly effective against *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Haemophilus influenzae*. The invention also relates to a process for the preparation of the compounds of the present invention, pharmaceutical compositions containing the compounds of the present invention and the methods for treating bacterial infection.

BACKGROUND OF THE INVENTION

The first generation macrolides erythromycin A and the early derivatives are characterized by bacteriostatic or bacteriocidal activity for most gram-positive bacteria, atypical pathogens, many communities acquired respiratory infections, in-patients with penicillin allergy. However, erythromycin A causes numerous drug-drug interactions, has relatively poor absorption, poor local tolerance, loses its antibacterial activity under acidic conditions by degradation and the degraded products are known to be responsible for undesired side effects. (Itoh, Z., et al., *Am. J. Physiol.*, (1984), 247: 688; and Omura, S., et al., *J. Med. Chem.*, (1987), 30, 1943). Various erythromycin A derivatives have been prepared to overcome the acid instability and other problems associated with it.

Roxithromycin, clarithromycin (6-O-methylerythromycin A) and azithromycin (azalides) have been developed to address the limitation of erythromycin A. Both clarithromycin and azithromycin have proved to be important drugs in the treatment and prophylaxis of atypical mycobacterial infectious in-patient with HIV.

Macrolides have proved to be effective drugs in the treatment of many respiratory tract infections. However, increasing resistance among *S. pneumoniae* has prompted the search for new compounds that retain the favourable safety profile, and a spectrum of activity confined to respiratory pathogens (*Expert Opinion Investigations Drugs*, (2001), 10, 353-367). Consequently, numerous investigators have prepared chemical derivatives

of erythromycin A in an attempt to obtain analogs having modified or improved profiles of antibiotic activity. It is reported that 6-O-methyl erythromycin A derivatives have improved acid stability and have superior *in vivo* antibacterial activity in comparison with erythromycin A when administered orally (Morimoto, et al., *Antibiotic*, (1984), 37, 187; *J. Antibiotic*, (1990), 43, 286; EP Pat. Nos. 272,110, and 215,355; and U.S. Pat. No. 4,331,803). It is reported that 6-O-methyl erythromycin derivatives and their 11, 12-cyclic carbamate derivatives have a superior *in vivo* antibacterial activity as well as stability to acids (U.S. Pat. No. 4,742,049; and EP Pat. No. 487,411). 6-O-methyl-3-descladinose erythromycin A (as in WO 97/10251), 6-O-substituted-3-oxoerythromycin A (as in U.S. Pat. No. 5,444,051), 3-deoxy-3-descladinose erythromycin A (as in WO 97/17356), tricyclic erythromycin A (as in WO 92/09614) and bicyclic 6-O-methyl 3-oxoerythromycin A (as in EP. Pat. No. 596,802) derivatives have been reported. There are also reports relating to ester derivatives at the C-3 position (EP. Pat. No. 619,320) with an enhanced spectrum of antibacterial activity. Ketolides exhibits greater efficacy and safety, has broader spectrum of activities, and is particularly effective against resistant pathogens, have been developed as next generation macrolides. 3'-N-modified-6-O-substituted erythromycin ketolides have been described (U.S. Pat. No. 6,034,69). Telithromycin (Aventis) and ABT-773 (Abbott) are expected to receive the FDA approval. They appeared to represent a useful alternative to macrolides and quinolones for treatment of community acquired respiratory infections.

Despite these advances, a need for better profile of antibiotic activity remains. Ideally, such derivatives would be water soluble with oral efficacy. Such compounds would provide useful agents for bacterially infectious diseases. There is no example available in the prior art wherein the compounds of the present invention, containing 3'-N-substituted-3-O-substituted erythronolide A derivatives are useful as therapy for curing bacterial infections.

SUMMARY OF THE INVENTION


Novel 3'-N-substituted-3-O-substituted erythronolide A derivatives are provided, which are useful for the safe treatment of bacterially infectious disease, and methods for the syntheses of these compounds are also provided. The pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers,

polymorphs of these compounds, as well as metabolites having same type of activity are also provided. Pharmaceutical compositions containing such compounds their pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, together with acceptable carriers, excipients or diluents, which are useful for the treatment of bacterially infectious disease are also provided.

The compounds disclosed herein were screened for antibacterial activity *in vitro* using agar incorporation method. Several compounds exhibited significant antibacterial activity. The 3'-N-substituted-3-O-substituted erythronolide A derivatives are useful as therapy for curing bacterial infections.

Other aspects will be set forth in the description, which follows and in part will be apparent from the description or may be learnt by the practice of the invention. Advantages which derive from the compounds and methods disclosed herein may be realized and obtained by means of the mechanism and combination pointed out in the appended claims.

In one aspect, there are provided compounds having structures of Formula I:



Formula I

and their pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein:

R¹ represents: lower alkyl (C₁-C₅) group, lower alkyl (C₁-C₅) having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C₁-C₅) amino group, lower alkyl amino (C₁-C₅) carbonyl group; lower alkoxy group (C₁-C₅); or five or six membered

aryl or heteroaryl ring having 1 to 3 hetero atoms such as oxygen, nitrogen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such as lower alkyl (C₁-C₅) group, lower alkyl (C₁-C₅) group having one or more halogen atoms, lower alkoxy (C₁-C₅) groups, lower alkyl (C₁-C₅) amino group, halogen atoms, amino group, nitro group, hydroxy group, and cyano group;

R² and R³ are independently selected from: C₁-C₆ alkyl group optionally substituted with halogen atoms; cycloalkyl (C₃-C₇) group; or five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom such as nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such as lower alkyl (C₁-C₃), lower alkyl (C₁-C₃) group having one or more halogen atom as substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, and cyano group; the above-mentioned C₁-C₆ alkyl group may be substituted by: NHCOR⁵, NHCOOR⁵, OCOR⁵, or COR⁵ wherein R⁵ represents lower alkyl (C₁-C₅); five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom such as nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such as lower alkyl (C₁-C₃), lower alkyl (C₁-C₃) group having one or more halogen atoms as substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino group, halogen atoms, nitro group, hydroxy group, and cyano group; C₂-C₆ alkenyl or alkyne group optionally substituted with halogen atoms or a group such as NHCOR⁵, NHCOOR⁵, COR⁵, or OCOR⁵ (wherein R⁵ is as defined above); cycloalkyl (C₃-C₇) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl (C₁-C₃) group, lower alkyl (C₁-C₃) group having one or more halogen atoms as substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group;

R' represents hydrogen, or a hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxy methyl;

R'' represents hydrogen, or a lower alkyl (C₁-C₃) group;

Y represents oxygen or sulphur;

Z represents an oxygen atom or a group represented by NOR^6 , wherein R^6 represents hydrogen atom, alkyl ($\text{C}_1\text{-C}_6$) group, alkyl ($\text{C}_1\text{-C}_6$) amino group, phenyl or benzyl group, or phenyl or benzyl group having 1 to 5 substituents such as halogen atoms, lower alkyl ($\text{C}_1\text{-C}_3$) group, hydroxy group, nitro group, cyano group, or amino group;

5 U represents a hydroxy group, OR^7 , wherein R^7 represents hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxymethyl; or $-\text{NH}(\text{CH}_2)_n\text{R}^8$, wherein n represents 0 to 4 and R^8 represents five or six membered aryl or heteroaryl ring having 1 to 4 hetero atom such as nitrogen, oxygen, and sulphur, wherein
 10 the aryl or heteroaryl ring may be unsubstituted or substituted by one to three substituents such as lower alkyl ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen atoms as substituent(s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino, halogen atoms, nitro group, hydroxy group, amino group, and cyano group;

V represents: hydrogen atom; hydroxy group; or OR^7 , wherein R^7 represents a hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl,
 15 methylthiomethylmethyl and methoxymethyl;

U and V may also together represent (with carbon atoms at the 11- and 12-positions on the erythronolide skeleton): a group represented by Formula



or a group represented by the Formula



wherein R^9 represents: hydrogen atom; alkyl ($\text{C}_1\text{-C}_6$) group, wherein the alkyl ($\text{C}_1\text{-C}_6$) may be unsubstituted or substituted by halogen atoms, five or six membered aryl or heteroaryl ring having 1 to 3 hetero atoms such as nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such
 30 as lower alkyl ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen (F, Cl, Br, I) atoms as substituent(s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group.

In a second aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection comprising administering to a patient in need thereof, a therapeutically effective amount of a compound as described above.

5 In a third aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection mediated through *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, or *Haemophilus influenzae*. with a compound as described above.

10 In a fourth aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection, comprising administering to a patient in need thereof, a therapeutically effective amount of a pharmaceutical composition including a compound as described above.

15 In a fifth aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection mediated through *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, or *Haemophilus influenzae* with a therapeutically effective amount of a pharmaceutical composition as described above.


In a sixth aspect, there are provided processes for preparing 3'-N-substituted-3-O-substituted erythronolide A derivatives.

20 The compounds described herein exhibit significant potency in terms of their antibacterial activity *in vitro*. Therefore, pharmaceutical compositions and methods for treatment of bacterially infectious diseases are provided herein. In addition, the compounds or pharmaceutical compositions disclosed herein can be administered orally or parentally.

25

DETAILED DESCRIPTION OF THE INVENTION

The compounds described herein may be prepared by the following reaction sequence.

Scheme 1

In accordance with Scheme 1, clarithromycin of Formula II is converted into particular desired compounds of Formula I by a series of steps as follows:

Step (1): Clarithromycin of Formula II is suspended in aqueous alcohol such as methanol, ethanol, propanol, or isopropanol, and is hydrolyzed with a mineral or organic acid such as hydrochloric acid or dichloroacetic acid at an ambient temperature for about 1 to 25 hours to give a compound of Formula III.

Step (2): The compound of Formula III is reacted with R'_2O or $R'X$ (where R' is a hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxymethyl, and X is a halogen atom) in the presence of an inorganic base such as sodium hydrogen carbonate, or potassium carbonate or an organic base such as triethylamine, pyridine, tributylamine, or 4-N-dimethylaminopyridine, in an inert solvent such as dichloromethane, dichloroethane, acetone, ethyl acetate, or tetrahydrofuran, at a temperature of from 0°C to 30°C , to give a compound of Formula IV..

Step (3): The compound of Formula IV is reacted with $R^1\text{COOH}$, $R^1\text{COX}$, $(R^1\text{CO})_2\text{O}$ or $R^1\text{COOR}^4$ (where R^1 is as defined for Formula I and R^4 is a group such as pivaloyl group, p-toluenesulfonyl group, isobutoxycarbonyl group, ethoxycarbonyl group or isopropoxycarbonyl group) and activating reagent such as dicyclohexylcarbodiimide (DCC) or 1-ethyl-3(3-dimethylaminopropyl) carbodiimide hydrochloride (EDCI) in the presence of an inorganic base such as sodium hydrogen carbonate, or potassium carbonate, or an organic base such as triethylamine, pyridine, tributylamine, or 4-dimethylaminopyridine, in an inert solvent such as dichloromethane, dichloroethane, acetone, ethyl acetate, or tetrahydrofuran, at a temperature of from 0°C to 30°C to give a compound of Formula V.


Step (4): The compound of Formula V is heated with alcohol as used in step (1), at a temperature of from 30°C to 100°C for about 5 to 25 hours to remove the protecting group (R') at the 2'-position of desosaminyl moiety, to obtain a compound of Formula VI.

Step (5): The compound of Formula VI is desmethylated at 3'-N-dimethyl group with N-iodosuccinamide and acetonitrile or iodine in the presence of a base such as sodium acetate followed by quench with sodium thiosulfonate to give a compound of

Formula VII. Alternatively, the reaction is carried out using reagents such as benzylchloroformate, allylchloroformate, or vinylchloroformate.


Step (6): The compound of Formula VII is reacted with a reagent represented by the Formula R^2CHO (or its precursor), R^2_2CO or R^2X (wherein R^2 is as defined for Formula I and X represents halogen atom) in the presence of a reducing agent such as sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride or in the presence of palladium and carbon catalyst in a protic or non-protic solvent such as hexane, toluene, methylene chloride, ethylene chloride, chloroform, tetrahydrofuran, N-methylpyrrolidinone, diethyl ether, bis-methoxymethyl ether, dimethylformamide, acetonitrile, acetone, or ethyl acetate, under hydrogen atmosphere to give a particular compound of Formula I (wherein $R^3=R''=CH_3$, $R'=H$, $U=V=OH$, and $Y=Z=O$).

The order of steps presented in Scheme I is not critical, and reactions in other orders can also be carried out to produce particular compounds of Formula I. For example, step (5) of Scheme I, desmethylation at the 3'-N-dimethyl group, can be carried out as a first step on clarithromycin of Formula II, to produce an N-desmethylated compound of Formula VIII




Formula VIII

with reagents and conditions as described above for Scheme I, step (5). This can be followed by introduction of R^2 (step (6) in Scheme I), to produce a compound of Formula IX


**Formula IX**

using reagents and conditions as described above for Scheme I, step (6). This can be then followed by hydrolysis of a compound of Formula IX to produce a compound of Formula X

**Formula X**


5

using reagents and conditions as described above for Scheme I, step (1). Reaction of a compound of Formula X with a reagent R'_2O or $R'X$ (R' and X as previously described) can then result in a compound of Formula XI using reagents and conditions as described above for Scheme I, step (2).

**Formula XI**

10

Reaction of a compound of Formula XI with reagent R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 (R^1 , X and R^4 as described above) can produce a compound of Formula XII using reagents and conditions as described above for Scheme I, step (3).




5

Formula XII

Deprotection of a compound of Formula XII can produce a compound of Formula I, using reagents and conditions as described for Scheme I, step (4).


In another example, steps (1) through (3) can be carried out on clarithromycin of
 10 Formula II to produce compound of Formula V. From this point, desmethylation of the compound of Formula V can produce a compound of Formula XIII



Formula XIII

using reagents and conditions as described above for Scheme I, step (5). Deprotection of a compound of Formula XIII can produce a compound of Formula VII, using reagents and
 15 conditions as described for Scheme I, step (4). Then, introduction of R^2 , as in Scheme I, step (6), can produce a compound of Formula I, using the described reagents and conditions.

In yet another example, steps (1) and (2) can be carried out on clarithromycin of Formula II to produce a compound of Formula IV. From this point, desmethylation of the compound of Formula IV can be carried out to produce a compound of Formula XIV.



5

Formula XIV

Introduction of R^2 , as in Scheme I, step (6), can produce compound of Formula XI, using the described reagents and conditions. Reaction of a compound of Formula XI with reagent R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 (R^1 , X and R^4 as described above) can produce a compound of Formula XII using reagents and conditions as described above for Scheme I, step (3). Deprotection of a compound of Formula XII can produce a compound of Formula I, using reagents and conditions as described for Scheme I, step (4).

10

Other feasible combinations of steps will be recognized by those of ordinary skill in the art, using reagents described herein, or appropriate alternative reagents known in the art for these transformations.

15


Preferred compounds according to the invention and capable of being produced by Scheme 1 include:

Compound No.	Chemical Name
5 1.	3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 1)
2.	3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 2)
10 3.	3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 3)
15 4.	3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 4)
5.	3-O-(2-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 5)
20 6.	3-O-(3-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 6)
7.	3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 7)
25 8.	3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 8)
9.	3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 9)
30 10.	3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 10)
35 11.	3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 11)
12.	3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 12)
40 13.	3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 13)
45 14.	3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminyl-6-O-methyl erythronolide (Compound No. 14)

15. 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-cyclopropyl] desosaminyl-6-O-methyl erythronolide A (Compound No. 15)
- 5 16. 3-O-(3-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-cyclopropyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 16)
17. 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl] desosaminyl-6-O-methyl erythronolide A (Compound No. 17)
- 10 18. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 18)
19. 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 19)
- 15 20. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 20)
21. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 21)
- 20 22. 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 22)
23. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 23)
- 25 24. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-(4-nitro) benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 24)
- 30 25. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 25)
26. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 26)
- 35 27. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 27)
28. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 28)
- 40 29. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 29)
- 45 30. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 30)

31. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 31)
- 5 32. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 32)
33. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 33)
- 10 34. 3-O-(2-Nitrophenyl) acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl) desosaminy-6-O-methyl erythronolide A (Compound No. 34)
35. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl erythronolide A (Compound No. 35)
- 15 36. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-isopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 36)
37. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 37)
- 20 38. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 38)
39. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 39)
40. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 40)
- 30 41. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminy-6-O-methyl erythronolide A (Compound No. 41)
42. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 42)
- 35 43. 3-O-(2-Pyridyl)acetyl-5-O-[3'-N-desmethyl-3'-N-benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 43)
44. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 44)
- 40 45. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 45)
- 45 46. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl erythronolide A (Compound No. 46)

47. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl erythronolide A (Compound No. 47)
- 5 48. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl erythronolide A (Compound No. 48)
49. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 49)
- 10 50. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 50)
51. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 51)
- 15 52. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 52)
53. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 53)
- 20 54. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 54)
55. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 55)
- 25 56. 3-O-(Phenyl)acetyl-5-O-[(3'-N-desmethyl-3'-N-cyclopropylmethyl)]desosaminy-6-O-methyl erythronolide A (Compound No. 56)
- 30 57. 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-(4-fluoro)benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 57)
58. 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 58)
- 35 59. 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 59)
60. 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 60)
- 40 61. 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 61)
- 45 62. 3-O-(2-Thiophene)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 62)

63. 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminy-6-O-methyl erythronolide A (Compound No. 63)
- 5 64. 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 64)
65. 3-O-(4-chlorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 65)
- 10 66. 3-O-(2-Methylphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 66)
67. 3-O-(2-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 67)
- 15 68. 3-O-(4-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 68)
69. 3-O-(4-Methoxyphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 69)
- 20 70. 3-O-(4-Methoxyphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 70)
- 25 71. 3-O-(1-Naphthyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 71)
72. 3-O-(1-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 72)
- 30 73. 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 73)
- 

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 April 2005 (07.04.2005)

PCT

(10) International Publication Number
WO 2005/030786 A1

(51) International Patent Classification⁷: C07H 17/08,
A61K 31/7048

(21) International Application Number:
PCT/IB2003/004166

(22) International Filing Date:
25 September 2003 (25.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): RAN-
BAXY LABORATORIES LIMITED [IN/IN]; 19, Nehru
Place, New Delhi, Delhi 110 019 (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DAS, Biswajit
[IN/IN]; 2nd Floor, J7/13 DLF II, Gurgaon, Haryana 122
001 (IN). SALMAN, Mohammad [IN/IN]; P - 2/19,
DLF City, Phase - II, Gurgaon, Haryana 122 001 (IN).
SHELKE, Sachin, Vilasrao [IN/IN]; Gautam Public
School, Kolpewadi, Tal-kopargaon, Ahmadnagar, Maha-
rastra 423 601 (IN). HAJARE, Atul, Kashinath [IN/IN];
House No. 700, Sector 17A, Gurgaon, Haryana, 122 001
(IN). TALUKDAR, Sanjay [IN/IN]; 64/3, Ashokgarh,
Kolkata, West Bengal 700 035 (IN). RATHY, Sujata

[IN/IN]; WZ-325 Rishi Nagar, Rani Bagh, Delhi, Delhi
110 034 (IN). PAL, Arani [IN/IN]; Basundhara Apartment
6E, 19/922, Raja S.C. Mullick Road, Kolkata, West Bengal
700 086 (IN). RATTAN, Ashok [IN/IN]; B-481 Sarita
Vihar, New Delhi, Delhi 110 044 (IN).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 3'-N-SUBSTITUTED-3-O-SUBSTITUTED ERYTHRONOLIDE A DERIVATIVES

(57) Abstract: This invention generally relates to macrolides, more particularly, the invention relates to 3-N-substituted-3-O-substituted erythronolide A derivatives, which are antibacterial agents effective against gram positive or gram negative bacteria and atypical pathogens. The compounds of this invention are more particularly effective against *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Haemophilus influenzae*. The invention also relates to a process for the preparation of the compounds of the present invention, pharmaceutical compositions containing the compounds of the present invention and the methods for treating bacterial infection.

WO 2005/030786 A1

3'-N-SUBSTITUTED-3-O-SUBSTITUTED ERYTHRONOLIDE A DERIVATIVES**FIELD OF INVENTION**

This invention generally relates to macrolides, more particularly, the invention relates to 3'-N-substituted-3-O-substituted erythronolide A derivatives, which are antibacterial agents effective against gram positive or gram negative bacteria and atypical pathogens. The compounds of this invention are more particularly effective against *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Haemophilus influenzae*. The invention also relates to a process for the preparation of the compounds of the present invention, pharmaceutical compositions containing the compounds of the present invention and the methods for treating bacterial infection.

BACKGROUND OF THE INVENTION

The first generation macrolides erythromycin A and the early derivatives are characterized by bacteriostatic or bacteriocidal activity for most gram-positive bacteria, atypical pathogens, many communities acquired respiratory infections, in-patients with penicillin allergy. However, erythromycin A causes numerous drug-drug interactions, has relatively poor absorption, poor local tolerance, loses its antibacterial activity under acidic conditions by degradation and the degraded products are known to be responsible for undesired side effects. (Itoh, Z., et al., *Am. J. Physiol.*, (1984), 247: 688; and Omura, S., et al., *J. Med. Chem.*, (1987), 30, 1943). Various erythromycin A derivatives have been prepared to overcome the acid instability and other problems associated with it.

Roxithromycin, clarithromycin (6-O-methylethromycin A) and azithromycin (azalides) have been developed to address the limitation of erythromycin A. Both clarithromycin and azithromycin have proved to be important drugs in the treatment and prophylaxis of atypical mycobacterial infectious in-patient with HIV.

Macrolides have proved to be effective drugs in the treatment of many respiratory tract infections. However, increasing resistance among *S. pneumoniae* has prompted the search for new compounds that retain the favourable safety profile, and a spectrum of activity confined to respiratory pathogens (*Expert Opinion Investigations Drugs*, (2001), 10, 353-367). Consequently, numerous investigators have prepared chemical derivatives

of erythromycin A in an attempt to obtain analogs having modified or improved profiles of antibiotic activity. It is reported that 6-O-methyl erythromycin A derivatives have improved acid stability and have superior *in vivo* antibacterial activity in comparison with erythromycin A when administered orally (Morimoto, et al., *Antibiotic*, (1984), 37, 187; *J. Antibiotic*, (1990), 43, 286; EP Pat. Nos. 272,110, and 215,355; and U.S. Pat. No. 4,331,803). It is reported that 6-O-methyl erythromycin derivatives and their 11, 12-cyclic carbamate derivatives have a superior *in vivo* antibacterial activity as well as stability to acids (U.S. Pat. No. 4,742,049; and EP Pat. No. 487,411). 6-O-methyl-3-descladinose erythromycin A (as in WO 97/10251), 6-O-substituted-3-oxoerythromycin A (as in U.S. Pat. No. 5,444,051), 3-deoxy-3-descladinose erythromycin A (as in WO 97/17356), tricyclic erythromycin A (as in WO 92/09614) and bicyclic 6-O-methyl 3-oxoerythromycin A (as in EP. Pat. No. 596,802) derivatives have been reported. There are also reports relating to ester derivatives at the C-3 position (EP. Pat. No. 619,320) with an enhanced spectrum of antibacterial activity. Ketolides exhibits greater efficacy and safety, has broader spectrum of activities, and is particularly effective against resistant pathogens, have been developed as next generation macrolides. 3'-N-modified-6-O-substituted erythromycin ketolides have been described (U.S. Pat. No. 6,034,69). Telithromycin (Aventis) and ABT-773 (Abbott) are expected to receive the FDA approval. They appeared to represent a useful alternative to macrolides and quinolones for treatment of community acquired respiratory infections.

Despite these advances, a need for better profile of antibiotic activity remains. Ideally, such derivatives would be water soluble with oral efficacy. Such compounds would provide useful agents for bacterially infectious diseases. There is no example available in the prior art wherein the compounds of the present invention, containing 3'-N-substituted-3-O-substituted erythronolide A derivatives are useful as therapy for curing bacterial infections.

SUMMARY OF THE INVENTION


Novel 3'-N-substituted-3-O-substituted erythronolide A derivatives are provided, which are useful for the safe treatment of bacterially infectious disease, and methods for the syntheses of these compounds are also provided. The pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers,

polymorphs of these compounds, as well as metabolites having same type of activity are also provided. Pharmaceutical compositions containing such compounds their pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, together with acceptable carriers, excipients or diluents, which are useful for the treatment of bacterially infectious disease are also provided.

The compounds disclosed herein were screened for antibacterial activity *in vitro* using agar incorporation method. Several compounds exhibited significant antibacterial activity. The 3'-N-substituted-3-O-substituted erythronolide A derivatives are useful as therapy for curing bacterial infections.

Other aspects will be set forth in the description, which follows and in part will be apparent from the description or may be learnt by the practice of the invention. Advantages which derive from the compounds and methods disclosed herein may be realized and obtained by means of the mechanism and combination pointed out in the appended claims.

In one aspect, there are provided compounds having structures of Formula I:



Formula I

and their pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein:

R¹ represents: lower alkyl (C₁-C₅) group, lower alkyl (C₁-C₅) having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C₁-C₅) amino group, lower alkyl amino (C₁-C₅) carbonyl group; lower alkoxy group (C₁-C₅); or five or six membered

aryl or heteroaryl ring having 1 to 3 hetero atoms such as oxygen, nitrogen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such as lower alkyl (C₁-C₅) group, lower alkyl (C₁-C₅) group having one or more halogen atoms, lower alkoxy (C₁-C₅) groups, lower alkyl (C₁-C₅) amino group, halogen atoms, amino group, nitro group, hydroxy group, and cyano group;

R² and R³ are independently selected from: C₁-C₆ alkyl group optionally substituted with halogen atoms; cycloalkyl (C₃-C₇) group; or five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom such as nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such as lower alkyl (C₁-C₃), lower alkyl (C₁-C₃) group having one or more halogen atom as substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, and cyano group; the above-mentioned C₁-C₆ alkyl group may be substituted by: NHCOR⁵, NHCOOR⁵, OCOR⁵, or COR⁵ wherein R⁵ represents lower alkyl (C₁-C₅); five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom such as nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such as lower alkyl (C₁-C₃), lower alkyl (C₁-C₃) group having one or more halogen atoms as substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino group, halogen atoms, nitro group, hydroxy group, and cyano group; C₂-C₆ alkenyl or alkyne group optionally substituted with halogen atoms or a group such as NHCOR⁵, NHCOOR⁵, COR⁵, or OCOR⁵ (wherein R⁵ is as defined above); cycloalkyl (C₃-C₇) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl (C₁-C₃) group, lower alkyl (C₁-C₃) group having one or more halogen atoms as substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group;

R' represents hydrogen, or a hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxy methyl;

R'' represents hydrogen, or a lower alkyl (C₁-C₃) group;

Y represents oxygen or sulphur;

Z represents an oxygen atom or a group represented by NOR^6 , wherein R^6 represents hydrogen atom, alkyl ($\text{C}_1\text{-C}_6$) group, alkyl ($\text{C}_1\text{-C}_6$) amino group, phenyl or benzyl group, or phenyl or benzyl group having 1 to 5 substituents such as halogen atoms, lower alkyl ($\text{C}_1\text{-C}_3$) group, hydroxy group, nitro group, cyano group, or amino group;

5 U represents a hydroxy group, OR^7 , wherein R^7 represents hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxymethyl; or $-\text{NH}(\text{CH}_2)_n\text{R}^8$, wherein n represents 0 to 4 and R^8 represents five or six membered aryl or heteroaryl ring having 1 to 4 hetero atom such as nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by one to three substituents
 10 such as lower alkyl ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen atoms as substituent(s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino, halogen atoms, nitro group, hydroxy group, amino group, and cyano group;

V represents: hydrogen atom; hydroxy group; or OR^7 , wherein R^7 represents a hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl,
 15 methylthiomethylmethyl and methoxymethyl;

U and V may also together represent (with carbon atoms at the 11- and 12- positions on the erythronolide skeleton): a group represented by Formula



or a group represented by the Formula



wherein R^9 represents: hydrogen atom; alkyl ($\text{C}_1\text{-C}_6$) group, wherein the alkyl ($\text{C}_1\text{-C}_6$) may be unsubstituted or substituted by halogen atoms, five or six membered aryl or heteroaryl ring having 1 to 3 hetero atoms such as nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents such
 30 as lower alkyl ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen (F, Cl, Br, I) atoms as substituent(s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group.

In a second aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection comprising administering to a patient in need thereof, a therapeutically effective amount of a compound as described above.

In a third aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection mediated through *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, or *Haemophilus influenzae* with a compound as described above.

In a fourth aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection, comprising administering to a patient in need thereof, a therapeutically effective amount of a pharmaceutical composition including a compound as described above.


In a fifth aspect, there is provided a method of treating or preventing an animal or human suffering from bacterial infection mediated through *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, or *Haemophilus influenzae* with a therapeutically effective amount of a pharmaceutical composition as described above.

In a sixth aspect, there are provided processes for preparing 3'-N-substituted-3-O-substituted erythronolide A derivatives.

The compounds described herein exhibit significant potency in terms of their antibacterial activity *in vitro*. Therefore, pharmaceutical compositions and methods for treatment of bacterially infectious diseases are provided herein. In addition, the compounds or pharmaceutical compositions disclosed herein can be administered orally or parentally.

DETAILED DESCRIPTION OF THE INVENTION

The compounds described herein may be prepared by the following reaction sequence.

Scheme 1

In accordance with Scheme 1, clarithromycin of Formula II is converted into particular desired compounds of Formula I by a series of steps as follows:

Step (1): Clarithromycin of Formula II is suspended in aqueous alcohol such as methanol, ethanol, propanol, or isopropanol, and is hydrolyzed with a mineral or organic acid such as hydrochloric acid or dichloroacetic acid at an ambient temperature for about 1 to 25 hours to give a compound of Formula III.

Step (2): The compound of Formula III is reacted with R'_2O or $R'X$ (where R' is a hydroxy protecting group such as acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxymethyl, and X is a halogen atom) in the presence of an inorganic base such as sodium hydrogen carbonate, or potassium carbonate or an organic base such as triethylamine, pyridine, tributylamine, or 4-N-dimethylaminopyridine, in an inert solvent such as dichloromethane, dichloroethane, acetone, ethyl acetate, or tetrahydrofuran, at a temperature of from 0°C to 30°C , to give a compound of Formula IV..

Step (3): The compound of Formula IV is reacted with $R^1\text{COOH}$, $R^1\text{COX}$, $(R^1\text{CO})_2\text{O}$ or $R^1\text{COOR}^4$ (where R^1 is as defined for Formula I and R^4 is a group such as pivaloyl group, p-toluenesulfonyl group, isobutoxycarbonyl group, ethoxycarbonyl group or isopropoxycarbonyl group) and activating reagent such as dicyclohexylcarbodiimide (DCC) or 1-ethyl-3(3-dimethylaminopropyl) carbodiimide hydrochloride (EDCI) in the presence of an inorganic base such as sodium hydrogen carbonate, or potassium carbonate, or an organic base such as triethylamine, pyridine, tributylamine, or 4-dimethylaminopyridine, in an inert solvent such as dichloromethane, dichloroethane, acetone, ethyl acetate, or tetrahydrofuran, at a temperature of from 0°C to 30°C to give a compound of Formula V.


Step (4): The compound of Formula V is heated with alcohol as used in step (1), at a temperature of from 30°C to 100°C for about 5 to 25 hours to remove the protecting group (R') at the 2'-position of desosaminyl moiety, to obtain a compound of Formula VI.

Step (5): The compound of Formula VI is desmethylated at 3'-N-dimethyl group with N-iodosuccinamide and acetonitrile or iodine in the presence of a base such as sodium acetate followed by quench with sodium thiosulfonate to give a compound of

Formula VII. Alternatively, the reaction is carried out using reagents such as benzylchloroformate, allylchloroformate, or vinylchloroformate.


Step (6): The compound of Formula VII is reacted with a reagent represented by the Formula R^2CHO (or its precursor), R^2_2CO or R^2X (wherein R^2 is as defined for Formula I and X represents halogen atom) in the presence of a reducing agent such as sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride or in the presence of palladium and carbon catalyst in a protic or non-protic solvent such as hexane, toluene, methylene chloride, ethylene chloride, chloroform, tetrahydrofuran, N-methylpyrrolidinone, diethyl ether, bis-methoxymethyl ether, dimethylformamide, acetonitrile, acetone, or ethyl acetate, under hydrogen atmosphere to give a particular compound of Formula I (wherein $R^3=R''=CH_3$, $R'=H$, $U=V=OH$, and $Y=Z=O$).

The order of steps presented in Scheme I is not critical, and reactions in other orders can also be carried out to produce particular compounds of Formula I. For example, step (5) of Scheme I, desmethylation at the 3'-N-dimethyl group, can be carried out as a first step on clarithromycin of Formula II, to produce an N-desmethylated compound of Formula VIII




Formula VIII

with reagents and conditions as described above for Scheme I, step (5). This can be followed by introduction of R^2 (step (6) in Scheme I), to produce a compound of Formula IX


**Formula IX**

using reagents and conditions as described above for Scheme I, step (6). This can be then followed by hydrolysis of a compound of Formula IX to produce a compound of Formula X

**Formula X**


5

using reagents and conditions as described above for Scheme I, step (1). Reaction of a compound of Formula X with a reagent R'_2O or $R'X$ (R' and X as previously described) can then result in a compound of Formula XI using reagents and conditions as described above for Scheme I, step (2).

**Formula XI**

10

Reaction of a compound of Formula XI with reagent $R^1\text{COOH}$, $R^1\text{COX}$, $(R^1\text{CO})_2\text{O}$ or $R^1\text{COOR}^4$ (R^1 , X and R^4 as described above) can produce a compound of Formula XII using reagents and conditions as described above for Scheme I, step (3).




5

Formula XII


Deprotection of a compound of Formula XII can produce a compound of Formula I, using reagents and conditions as described for Scheme I, step (4).

In another example, steps (1) through (3) can be carried out on clarithromycin of
 10 Formula II to produce compound of Formula V. From this point, desmethylation of the compound of Formula V can produce a compound of Formula XIII

**Formula XIII**

using reagents and conditions as described above for Scheme I, step (5). Deprotection of a compound of Formula XIII can produce a compound of Formula VII, using reagents and
 15 conditions as described for Scheme I, step (4). Then, introduction of R^2 , as in Scheme I, step (6), can produce a compound of Formula I, using the described reagents and conditions.

In yet another example, steps (1) and (2) can be carried out on clarithromycin of Formula II to produce a compound of Formula IV. From this point, desmethylation of the compound of Formula IV can be carried out to produce a compound of Formula XIV.



5

Formula XIV

Introduction of R^2 , as in Scheme I, step (6), can produce compound of Formula XI, using the described reagents and conditions. Reaction of a compound of Formula XI with reagent R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 (R^1 , X and R^4 as described above) can produce a compound of Formula XII using reagents and conditions as described above for Scheme I, step (3). Deprotection of a compound of Formula XII can produce a compound of Formula I, using reagents and conditions as described for Scheme I, step (4).

Other feasible combinations of steps will be recognized by those of ordinary skill in the art, using reagents described herein, or appropriate alternative reagents known in the art for these transformations.

15

Preferred compounds according to the invention and capable of being produced by Scheme 1 include:

Compound No.	Chemical Name
5	1. 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 1)
	2. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 2)
10	3. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 3)
15	4. 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 4)
	5. 3-O-(2-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 5)
20	6. 3-O-(3-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 6)
	7. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 7)
25	8. 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 8)
	9. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 9)
30	10. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 10)
35	11. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 11)
	12. 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 12)
40	13. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 13)
45	14. 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminyl-6-O-methyl erythronolide (Compound No. 14)

15. 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-cyclopropyl] desosaminyl-6-O-methyl erythronolide A (Compound No. 15)
- 5 16. 3-O-(3-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-cyclopropyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 16)
17. 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl] desosaminyl-6-O-methyl erythronolide A (Compound No. 17)
- 10 18. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 18)
19. 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 19)
- 15 20. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 20)
21. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 21)
- 20 22. 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 22)
23. 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 23)
- 25 24. 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-(4-nitro) benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 24)
- 30 25. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 25)
26. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 26)
- 35 27. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 27)
28. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 28)
- 40 29. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 29)
- 45 30. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 30)

31. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 31)
- 5 32. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 32)
33. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 33)
- 10 34. 3-O-(2-Nitrophenyl) acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl) desosaminy-6-O-methyl erythronolide A (Compound No. 34)
35. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl erythronolide A (Compound No. 35)
- 15 36. 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-isopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 36)
- 20 37. 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 37)
38. 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-methyl erythronolide A (Compound No. 38)
- 25 39. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 39)
40. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 40)
- 30 41. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl) desoaminy-6-O-methyl erythronolide A (Compound No. 41)
42. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 42)
- 35 43. 3-O-(2-Pyridyl)acetyl-5-O-[3'-N-desmethyl-3'-N-benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 43)
- 40 44. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 44)
- 45 45. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl erythronolide A (Compound No. 45)
46. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl erythronolide A (Compound No. 46)

47. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 47)
- 5 48. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 48)
49. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 49)
- 10 50. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 50)
51. 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 51)
- 15 52. 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 52)
53. 3-O-(4-Pyridyl) acetyl-5-O-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No.53)
- 20 54. 3-O-(4-Pyridyl) acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 54)
55. 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 55)
- 25 56. 3-O-(Phenyl)acetyl-5-O-[(3'-N-desmethyl-3'-N-cyclopropylmethyl]desoaminyl-6-O-methyl erythronolide A (Compound No. 56)
- 30 57. 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-(4-fluoro)benzyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 57)
58. 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 58)
- 35 59. 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 59)
60. 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 60)
- 40 61. 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 61)
- 45 62. 3-O-(2-Thiophene)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 62)

63. 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminy-6-O-methyl erythronolide A (Compound No. 63)
- 5 64. 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 64)
65. 3-O-(4-chlorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 65)
- 10 66. 3-O-(2-Methylphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 66)
67. 3-O-(2-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 67)
- 15 68. 3-O-(4-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 68)
69. 3-O-(4-Methoxyphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 69)
- 20 70. 3-O-(4-Methoxyphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 70)
- 25 71. 3-O-(1-Naphthyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 71)
72. 3-O-(1-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 72)
- 30 73. 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 73)
74. 3-O-(2,4-Difluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 74)
- 35 75. 3-O-(2,4-Difluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 75)
- 40 76. 3-O-(2-Bromophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 76)
77. 3-O-(2-Bromophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 77)
- 45 78. 3-O-(3-Indole)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminy-6-O-methyl erythronolide A (Compound No. 78)

79: 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl
erythronolide A. (Compound No. 79)

5 The compounds described herein can be used as antibacterial agents, effective
against gram positive or gram negative and atypical pathogens. The compounds are, in
particular, effective against *Staphylococcus aureus*, *Streptococcus pneumoniae*,
Enterococcus faecalis, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Haemophilus*
influenzae.

10 Functional groups contained in compounds described herein can form organic or
inorganic acid salts, which are within the scope of sound medical judgement suitable for
use in contact with the tissue of humans and lower animals without undue toxicity,
irritation, allergic response and the like. Also, other amino groups (primary and
secondary) can be present, and can also form organic or inorganic salts. The resulting salts
are useful by themselves and in the therapeutic composition. These salts may be prepared
15 by conventional techniques, such as suspending the compound in water and then adding
one equivalent of the organic acid such as acetic acid, oxalic acid, maleic acid, tartaric
acid, citric acid, succinic acid, malonic acid, adipic acid, ascorbic acid, camphorenic acid,
nicotinic acid, butyric acid, lactic acid, or glucuronic acid, or inorganic acids such as
hydrochloric acid, hydrobromic acid, phosphoric acid, sulphuric acid, nitric acid, boric
20 acid or perchloric acid.

The neutral solution of the resulting salt is subjected to rotary evaporation under
diminished pressure to the volume necessary to ensure precipitation of the salt upon
cooling, which is then filtered and dried. The salts of the compounds described herein
may also be prepared under strictly non-aqueous conditions. For example, dissolving free
25 base in a suitable organic solvent, adding one equivalent of the desired acid to the same
solvent and stirring the solution at 0-5 °C, causes the precipitation of the acid addition salt,
which is then filtered, washed the solvent, and dried.

Alternatively, the solvent is stripped off completely to obtain the desired salt.
These salts are often preferred for use in formulating the therapeutic compositions
30 described herein, because they are crystalline and relatively recently more stable and
water-soluble.

Making use of their antibacterial activity, the compounds disclosed herein may be administered to an animal for treatment orally, topically, rectally, intranasally, or by parenteral route. The pharmaceutical compositions described herein comprise a pharmaceutically effective amount of a compound disclosed herein formulated together with one or more pharmaceutically acceptable carriers. The term "pharmaceutically acceptable carriers" is intended to include non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

Solid form preparations for oral administrations include capsules, tablet, pills, powder, granules, cachets and suppositories. For solid form preparations, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier, such as sodium citrate, dicalcium phosphate and/or a filler or extenders such as starches, lactose, sucrose, glucose, mannitol or silicic acid; binders such as carboxymethylcellulose, alginates, gelatins, polyvinylpyrrolidinone, sucrose, or acacia; disintegrating agents such as agar-agar, calcium carbonate, potato starch, alginic acid, certain silicates or sodium carbonate; absorption accelerators such as quaternary ammonium compounds; wetting agents such as cetyl alcohol, glycerol mono stearate; adsorbants such as Kaolin; lubricants such as talc, calcium stearate, magnesium stearate, solid polyethyleneglycol, sodium lauryl sulphate or mixtures thereof. In the case of capsules, tablets, or pills, the dosage form may also comprise buffering agents.

The solid preparation of tablets, capsules, pills, or granules can be prepared with coating and shells such as enteric coating and other coatings well known in the pharmaceutical formulating art.

Liquid form preparations for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. For liquid form preparation, the active compound is mixed with water or other solvent, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (such as cottonseed, groundnut, corn, germ, olive, castor and sesame oil), glycerol, and fatty acid esters of sorbitan and mixture thereof. Besides inert diluents, the oral composition can also include adjuncts, such as wetting agents, emulsifying agents, suspending agents, sweetening agents, flavouring agents and perfuming agents.

Injectable preparations such as sterile injections, aqueous or oleaginous suspensions may be formulated according to the art using suitable dispersing or wetting and suspending agent. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U. S. P. and isotonic sodium chloride.

5 Dosage forms for tropical or transdermal administration of a compound of the present invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active compound is admixed under sterile condition with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulations, ear drops, eye ointments, powder and solution are
10 also contemplated as being within the scope of this invention.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be packaged preparation, the package containing discrete capsules, powders, in vials or ampules, and ointments capsule, cachet, tablet, gel,
15 cream itself or it can be the appropriate number of any of these packaged forms.

The quantity of active compound in unit dose of preparation may be varied or adjusted from less than 1 mg to several grams according to the particular application and potency of the active ingredient.

For therapeutic use, as agents for treating bacterial infections, the compounds
20 disclosed here are utilized at initial dosages of about 3 mg to about 40 mg per kilogram daily. The dosages, however, may be varied depending upon the requirements of the patients and the compound being employed.

Pharmaceutically acceptable acid addition salts, solvates, enantiomers, diastereomers, polymorphs and metabolites having the same type of physiological or or
25 therapeutic activity can also be used as therapeutic agents. Pharmaceutical compositions comprising the compounds disclosed herein, their enantiomers, diastereomers, pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, polymorphs, metabolites in combination with pharmaceutically acceptable carrier and optionally included excipients are also provided hereby.

In the above syntheses, where specific acids, bases, solvents, catalysts, reducing agents etc., are mentioned, it is to be understood that the other acids, bases, solvents, catalysts, reducing agents etc, may be used. Similarly, the reaction temperature and duration of the reaction may be adjusted according to the need.

5 The examples mentioned below demonstrate the general synthetic procedures as well as the specific preparation of particular compounds. The examples are provided to illustrate the details of the invention and should not be constrained to limit the scope of the present invention, which is defined by the claims appended hereto.

10 **EXPERIMENTAL DETAILS**

Various solvents, such as acetone, methanol, ethyl acetate, ether, tetrahydrofuran, pyridine, hexane and dichloromethane were dried using various drying agents according to procedures described in the literature. IR spectra were recorded as nujol mull or a thin neat film on a Perkin Elmer Paragone instrument, Nuclear Magnetic Resonance (NMR) were recorded on a Varion XL-300 MHz instrument using tetramethylsilane as an internal standard.

EXAMPLE 1: Preparation of 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl erythronolide A (Compound No. 56)

20 Step 1 : 5-O-Desosaminyl-6-O-methyl erythronolide A

To a solution of hydrochloric acid (~1N, 160 mL) was added clarithromycin (10 g, 13.37 mmol) in portion at ambient temperature. After addition, the reaction mixture was stirred at 20° to 40° C for about 40 minutes. The reaction mixture was neutralized with solid sodium bicarbonate (NaHCO₃) in cold and extracted with ethyl acetate. The organic layer was washed with water, brine, dried over anhydrous sodium sulphate (Na₂SO₄) and then the solvent was removed under vacuum to give white solid, 7.5 g (yield, 96%).

Step 2 : 3-Hydroxy-5-O-(2'-O-benzoyl) desosaminy-6-O-methyl erythronolide A

5-O-Desosaminyl-6-O-methyl erythronolide A (7 g, 11.86 mmol, from step 1) was taken in dichloromethane (120 mL) and to it was added triethylamine (6.58 mL) and benzoic anhydride (6.7 g, 29.61 mmol) under argon atmosphere at 20° to 40°C. The reaction mixture was stirred at ambient temperature under argon atmosphere for about 48 hours. The reaction mixture was diluted with dichloromethane and the organic layer was

washed successively with saturated sodium bicarbonate (NaHCO_3) solution and brine before drying over anhydrous sodium sulphate (Na_2SO_4). Removal of solvent afforded a white foamy solid, which was purified by column chromatography using 2% methanol in dichloromethane as eluent to obtain the desired product, 5.6 g (yield, 68 %).

5 **Step 3: 3-O-(Phenyl)acetyl-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A**

3-Hydroxy-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A (1 g, 1.44 mmol, from step 2) was dissolved in dry dichloromethane (30 mL), to this solution was added phenyl acetic acid (588 mg, 4.32 mmol), 1-ethyl-3-(3-dimethylaminopropyl)
10 carbodiimide hydrochloride (40 mg, 2.1 mmol), 4-dimethylaminopyridine (256 mg, 2.1 mmol) under ice cooling condition. The whole reaction mixture was stirred at 0 °C to ambient temperature for about 6 hours. This was poured into 5% sodium bicarbonate (NaHCO_3) solution and extracted with dichloromethane. Organic layer was washed with water, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under
15 vacuum. The crude product was then purified by column chromatography using hexane:acetone:triethylamine (9:1:0.1) as eluent to obtain the desired product, 700 mg (yield, 60 %).

The following compounds were prepared by using above-mentioned sequence, using the appropriate acetic acid in step 3.

- 20 3-O-(3-Nitrophenyl)acetyl-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A
3-O-(4-Nitrophenyl)acetyl-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A
3-O-(2-Nitrophenyl)acetyl-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A
3-O-(2-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl)-desosaminyl-6-O-methyl erythronolide A
3-O-(3-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl)-desosaminyl-6-O-methyl erythronolide A
25 3-O-(4-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl)-desosaminyl-6-O-methyl erythronolide A
3-O-(4-Chlorophenyl)acetyl-5-O-(2'-O-benzoyl)-desosaminyl-6-O-methyl erythronolide A
3-O-(2-Thiophene)acetyl-5-O-(2'-O-benzoyl)-desosaminyl-6-O-methyl erythronolide A
3-O-(4-Methoxyphenyl)acetyl-5-O-(2'-O-benzoyl)-desosaminyl-6-O-methyl erythronolide A
A

3-O-(4-Pyridyl)acetyl-5-O-(2'-O-benzoyl)-desosaminy-6-O-methyl erythronolide A

Step 4 : 3-O-(Phenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(Phenyl)acetyl-5-O-(2'-O-benzoyl)desosaminy-6-O-methyl erythronolide A (2.9 g, 3.57 mmol, from step 3) was taken in methanol (60 mL). The whole reaction mixture was heated at 60 °C for about 20 hours. Methanol was removed and the residue was purified by column chromatography using hexane: acetone: triethylamine (9:1:0.1) as eluent to obtain the desired product, 2.4 g (yield, 94 %).

The following compounds were prepared analogously.

3-O-(4-Nitrophenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

10 3-O-(3-Nitrophenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(2-Nitrophenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(2-Fluorophenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(3-Fluorophenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(4-Fluorophenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

15 3-O-(4-Chlorophenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(4-Methoxyphenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(2-Thiophene)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

3-O-(4-Pyridyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A

Step 5 : 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl)desosaminy-6-O-methyl erythronolide A

To a solution of 3-O-(Phenyl)acetyl-5-O-desosaminy-6-O-methyl erythronolide A (500 mg, 0.71 mmol, from step 4) in dry acetonitrile (20 mL), was added N-iodosuccinimide (179 mg, 0.79 mmol) in portion. The whole reaction mixture was stirred for about 15-20 hours. To this reaction mixture was added sodium bisulphite (NaHSO₃) solution till the colour become pale yellow. To this was added saturated sodium bicarbonate (NaHCO₃) solution. The whole reaction mixture was stirred for about 1 hour and then extracted with ethyl acetate. Organic layer was washed with water, dried over anhydrous sodium sulphate (Na₂SO₄) and the solvent was removed under vacuum. The crude product was then purified by column chromatography using hexane: acetone: TEA (1:1:0.1) to obtain the desired product, 450 mg (yield, 29%).

The following compounds were prepared analogously.

- 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 5 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 10 3-O-(4-Methoxyphenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 15 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 20 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- Step 6 : 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A**

3-O-(phenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl erythronolide A (200 mg, 0.29 mmol, from step 5) was dissolved in dry acetonitrile (5 mL), to this was added

25 anhydrous sodium bicarbonate (NaHCO_3 , 120 mg, 1.43 mmol), followed by bromomethylcyclopropane (75 mg, 0.5 mmol). The whole reaction mixture was stirred at ambient temperature under argon atmosphere for about 60 hours. The reaction mixture was diluted with ethyl acetate and the organic layer was washed with water, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum. This

30 crude (Na_2SO_4) product was then purified by column chromatography using hexane:acetone:triethylamine (95:5:1) as eluent to obtain the desired product, 120 mg (56%). $m/z=766$ (M+H)

The following compounds were prepared analogously.

3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 2)

$m/z = 766$ (M+H)

5 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 3)

$m/z = 766$ (M+H)

3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 25)

10 $m/z = 793$ (M+H)

3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 26)

$m/z = 793$ (M+H)

15 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 59)

$m/z = 754$ (M+H)

3-O-(4-Methoxyphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 69)

$m/z = 778$ (M+H)

20 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 50)

$m/z = 749$ (M+H)

3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 49)

25 $m/z = 749$ (M+H)

3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 31)

$m/z = 779$ (M+H)

30 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 33)

$m/z = 779$ (M+H)

3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 32)

$m/z = 779$ (M+H)

5

3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 40)

$m/z = 749$ (M+H)

10 EXAMPLE 2: Preparation of 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 4)

3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A (350 mg., 0.49 mmol, example 1, step 5) was dissolved in methanol, to this was added 4-fluorobenzaldehyde (121 mg, 0.91 mmol) and glacial acetic acid (294 mg, 4.9 mmol). The whole reaction mixture was stirred at 0° C for about 1.0 hours, then sodium cyanoborohydride (60 mg, 0.983 mmol) was added. The whole reaction mixture was stirred overnight. The reaction mixture was diluted with ethyl acetate followed by sodium bicarbonate (NaHCO₃) solution. Organic layer was washed with water, dried over anhydrous sodium sulphate and the solvent was removed under vacuum. This crude product was then purified by column chromatography using hexane:acetone: triethylamine (9:1:0.1) as eluent to obtain the desired product, 200 mg (yield, 50%).

$m/z = 820$ (M+H)

The following compounds were prepared analogously using the appropriate aldehyde.

25 3-O-(2-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 5)

$m/z = 820$ (M+H)

3-O-(3-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 6)

30 $m/z = 820$ (M+H)

3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 7)

$m/z = 752$ (M+H)

This title compound can be prepared using ethoxy trimethylsilyloxycyclopropyl instead of benzaldehyde.

5 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 27)

$m/z = 829$ (M+H)

3-O-(Phenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 57)

$m/z = 802$ (M+H)

10

EXAMPLE 3: Preparation of 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 8)

Step 1: 3'-N-Desmethyl Clarithromycin

To a solution of clarithromycin (10 g, 13.35 mmol) in dry acetonitrile (200 mL)
15 was added N-iodosuccinimide (4.2 g, 18.66 mmol) in portion. The whole reaction mixture was stirred at ambient temperature for about 24 hours. To this reaction mixture was added sodium bisulphite (NaHSO_3) and then sodium carbonate (Na_2CO_3) solution. The whole reaction mixture was then stirred for about 3 hours and extracted with ethyl acetate, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum.
20 The crude product was purified by column chromatography using hexane:acetone:triethylamine (6:1:0.1) as eluent to obtain the desired product, 9.3 g (yield, 47.4%).

Step 2: 3'-N-Desmethyl-3'-N-ethyl clarithromycin

Method A: To a solution of 3'-N-desmethyl clarithromycin (2 g, 2.7 mmol, from step 1) in dry dimethylformamide (DMF, 15 mL) was added anhydrous potassium
25 carbonate (K_2CO_3 , 1.80 g, 13.08 mmol). The whole reaction mixture was stirred for about 5 minutes. Then ethyl iodide (0.61 g, 3.91 mmol) was added. The whole reaction mixture was stirred at ambient temperature for about 14-16 hours. It was filtered through the celite bed. Dimethylformamide (DMF) was removed under vacuum. The residue was dissolved in ethyl acetate and washed with water. The organic layer was dried over sodium sulphate
30 (Na_2SO_4) and the solvent was removed under vacuum. The crude product was then purified by column chromatography using hexane:acetone:triethylamine (9:1:0.1) as eluent to obtain the desired product, 1 g (yield, 48%).

Method B: To a solution of 3'-N-desmethyl clarithromycin (2 g, 2.72 mmol, from step 1) in acetonitrile (30 mL) was added ethyl iodide (4.26 g, 27.32 mmol) and diisopropylethylamine (3.52 g, 27.32 mmol). The reaction mixture refluxed for about 4-5 hours. The reaction mixture was diluted with the dichloromethane and washed with 5% sodium bicarbonate (NaHCO_3) solution. The organic layer was dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum. It was purified by column chromatography using hexane:acetone:triethylamine (9:1:0.1) to obtain the desired product, 1.1 g (yield, 53%).

Step 3: 3-Hydroxy-5-O-(3'-N-desmethyl-3'-N-ethyl) desosaminyl-6-O-methyl erythronolide A

3'-N-Desmethyl-3'-N-ethyl-6-O-methyl erythronolide A (4.5 g, 5.89 mmol, from step 2) was added to a cold solution of hydrochloride (~1N, 60 mL). It was stirred for about 1 hour, neutralized with solid sodium bicarbonate (NaHCO_3) and extracted with ethyl acetate, dried over sodium sulphate (Na_2SO_4) and concentrated. The crude product was purified by column chromatography using 20 % acetone in hexane to obtain the desired product, 2.9 g (yield, 81%).

Step 4: 3-Hydroxy-5-O-(2'-benzoyl-3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl erythronolide A

3-Hydroxy-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl erythronolide A (2.9 g, 4.80 mmol, from step 3) was dissolved in dry dichloromethane. To this was added benzoic anhydride (2.15 g, 9.60 mmol) and triethylamine (2.2 g, 22 mmol). The whole reaction mixture was stirred at ambient temperature for about 12 hours. More benzoic anhydride (1.07 g, 2.4 mmol) was added, and reaction mixture was stirred for an additional 6 hours. The organic layer was washed with sodium bicarbonate (NaHCO_3) solution and then with water, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum to obtain crude product which on purification by column chromatography using 10 % acetone in hexane gave the desired product, 1.8 g (yield, 53%).

Step 5: 3-O-(3-Fluorophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

To a cold solution of 3-hydroxy-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A (400 mg, 0.56 mmol, from step 4) was added DCC (230 mg, 1.30 mmol), DMAP (139 mg, 1.31 mmol), pyridine (223 mg, 2.8 mmol) and 3-fluorophenyl acetic acid (174 mg, 1.31 mmol). The reaction mixture was stirred at 0° to 20 °C for about 12 hours. Another equivalent each of 3-fluorophenyl acetic acid and DCC were added. The reaction mixture was stirred for about 6 hours more, filtered through celite bed, washed sequentially with sodium bicarbonate (NaHCO₃) solution, dilute hydrochloride and finally with water. The organic layer was dried over anhydrous sodium sulphate (Na₂SO₄), and the solvent was removed under vacuum. The crude product was purified by column chromatography using hexane:acetone:triethylamine (9:1:0.1) to obtain the desired product, 400 mg (yield, 83.8%).

The following compounds were prepared analogously, using the appropriate acetic acid.

3-O-(2-Fluorophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(4-Fluorophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(2-Nitrophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(3-Nitrophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(4-Nitrophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(2-Pyridyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(Phenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(3-Indole)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(2-Naphthalene)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

3-O-(2-Thiophene)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

- 5 3-O-(4-Chlorophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A

Step 6: 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A

- 10 3-O-(3-Fluorophenyl)acetyl-5-O-[(2'-O-benzoyl)-3'-N-desmethyl-3'-N-ethyl]desoaminyl-6-O-methyl erythronolide A (350 mg, 0.42 mmol) was taken in methanol (60 mL) and refluxed for about 16 hours. Methanol was removed to get a gummy solid. This was purified by column chromatography using 2 % methanol in chloroform as eluent to obtain the desired product, 0.29 g (94.7 %).

- 15 $m/z = 740$ (M+H)

The following compounds were prepared analogously.

3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 52) $m/z = 723$ (M+H)

- 20 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 51) $m/z = 723$ (M+H)

3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 9) $m/z = 740$ (M+H)

- 25 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 10) $m/z = 740$ (M+H)

3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 28) $m/z = 767$ (M+H)

3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 29) $m/z = 767$ (M+H)

- 30 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 30) $m/z = 767$ (M+H)

3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 41) $m/z = 723$ (M+H)

3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A $m/z = 761$ (M+H) (Compound No. 80)

3-O-(3-Indole)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 78) $m/z = 761$ (M+H)

5 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A $m/z = 772$ (M+H) (Compound No. 79)

3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A $m/z = 728$ (M+H) (Compound No. 60)

10 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl erythronolide A $m/z = 762$ (M+H) (Compound No. 63)

EXAMPLE 4: Preparation of 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desoaminyl-6-O-methyl erythronolide A (Compound No. 58)

Step 1: 3-Hydroxy-5-O-desosaminyl-6-O-methyl erythronolide A

15 To a solution of hydrochloride (~1N, 160 mL) was added clarithromycin (10 g, 13.37 mmol) in portion over a period of about 15 minutes. The reaction mixture was neutralized with solid sodium bicarbonate (NaHCO_3) in cold and extracted with ethyl acetate. The organic layer was washed with water, brine, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed to give a crude product, 7.2 g (yield, 20 91%).

Step 2: 3-Hydroxy-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A

A solution of 3-hydroxy-5-O-desosaminyl-6-O-methyl erythronolide A (15 g, 25.4 mmol), benzoic anhydride (14.4 g, 64 mmol), triethylamine (12.8 g, 127 mmol) in dry dichloromethane (150 mL) was stirred at 30 °C for a period of about 48 hours. The organic 25 matter was extracted with dichloromethane, washed successively with sodium bicarbonate (NaHCO_3), water, brine, and dried over sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum to get crude residue. The crude product was purified by column chromatography over a SiO_2 bed thoroughly neutralized with triethyl amine (gradient elution with a 10-20% acetone in hexanes), 10.4 g (yield, 59%).

30 **Step 3: 3-O-(Phenyl)acetyl-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A**

To a solution of 3-hydroxy-5-O-(2'-benzoyl)desosaminyl-6-O-methyl erythronolide A (2.0 g, 2.9 mmol), DCC (1.45 g, 7.2 mmol), DMAP (0.9 g, 7.2 mmol), pyridine (1.1 g, 14.4 mmol) in dry dichloromethane (30 mL) was added phenyl acetic acid (0.98 g, 7.2 mmol) at 0 °C and the reaction mixture was allowed to attain 30 °C and stirred at that temperature for a period about 8-15 hours. The solid residue was removed by filtration over a celite bed. The organic matter was extracted with dichloromethane, washed successively with sodium bicarbonate (NaHCO₃), water, brine, and dried over anhydrous sodium sulphate (Na₂SO₄) and the solvent was removed under vacuum to obtain crude residue. The pure product was obtained by column chromatography over a SiO₂ bed thoroughly neutralized with triethylamine (gradient elution with a 10-20% acetone in hexanes), 1.7 g (74 %).

Step 4: 3-O-(Phenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A

To a solution of 3-O-(phenyl)acetyl-5-O-(2'-benzoyl)desosaminyl-6-O-methyl erythronolide A (1.7 g, 2.1 mmol) in dry acetonitrile (80 mL) was added N-iodosuccinimide (0.61g, 2.73 mmol) at 0 °C and the reaction mixture was allowed to attain 30 °C and stirred at that temperature for a period of more than 24 hours. The reaction mixture was stirred with a 5% solution of sodium bisulphite followed by stirring with 5% sodium carbonate solution. The organic matter was extracted with ethyl acetate, washed successively with water, brine, and dried over anhydrous sodium sulphate (Na₂SO₄) and the solvent was removed under vacuum to obtain crude residue. The pure product was obtained by column chromatography over a SiO₂ bed thoroughly neutralized with triethylamine (gradient elution with a 10-20% acetone in hexanes), 1.0 g (60%).

The following compounds were prepared analogously using the appropriate acetic acid in step 3.

3-O-(2-Thiophene)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A

3-O-(4-Chlorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A

3-O-(2-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A

3-O-(4-Chlorophenyl)acetyl-5-O-(2-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A

- 3-O-(3-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(3-Pyridyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 5 3-O-(4-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2-Naphthyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 10 3-O-(4-Methoxyphenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2,4-Difluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2,4-Difluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 15 3-O-(2-Bromophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2-Bromophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 20 3-O-(1-Naphthyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(1-Naphthyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(2-Methylphenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 25 3-O-(2-Methylphenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Methylphenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 30 3-O-(4-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A
- Step 5: 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A**

A solution of 3-O-(phenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A (0.5 g, 0.62 mmol) in methanol (30 mL) was heated at 75-80 °C for about 8 hours. Removal of the solvent under vacuum yielded a crude residue. Purification of the crude product over a bed of silica gel thoroughly neutralized with
5 triethylamine (gradient elution with a 20-30% acetone in hexanes) afforded the desired product, 0.28 g (66%).

The following compounds were prepared analogously.

- 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 10 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 15 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 20 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(4-Methoxyphenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 25 3-O-(2,4-Difluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(2,4-Difluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(2-Bromophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 30 3-O-(2-Bromophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A
- 3-O-(1-Naphthyl)acetyl-5-O-(3'-N-desmethyl)desosaminyll-6-O-methyl erythronolide A

3-O-(1-Naphthyl)acetyl-5-O-(3'-N-desmethyl)desosaminy-6-O-methyl erythronolide A

3-O-(2-Methylphenyl)acetyl-5-O-(3'-N-desmethyl)desosaminy-6-O-methyl erythronolide A

5 3-O-(2-Methylphenyl)acetyl-5-O-(3'-N-desmethyl)desosaminy-6-O-methyl erythronolide A

3-O-(4-Methylphenyl)acetyl-5-O-(3'-N-desmethyl)desosaminy-6-O-methyl erythronolide A

3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl)desosaminy-6-O-methyl erythronolide A

10 **Step 6: 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-phenylmethyl)desosaminy-6-O-methyl erythronolide A**

To a solution of 3-O-(phenyl)acetyl-5-O-(3'-N-desmethyl)desosaminy-6-O-methyl erythronolide A (0.5 g, 0.72 mmol) in methanol (10 mL) were added benzaldehyde (0.38 g, 3.6 mmol), acetic acid (0.43 g, 7.3 mmol), NaBH₃CN (0.23 g, 3.6 mmol) sequentially at 15 0 °C and the reaction mixture was then allowed to attain room temperature, and stirred for a period of about 6–8 hours. Removal of the solvent under vacuum yielded a crude residue. Purification of the crude product over a bed of silica gel thoroughly neutralized with triethylamine (gradient elution with a 2–5% acetone in hexanes) afforded the title compound, 0.15 g (26.5%) $m/z = 784$ (M+H)

20 The following compounds were prepared analogously, using the appropriate benzaldehyde or derivative.

3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 61) $m/z = 790$ (M+H)

25 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 64) $m/z = 718$ (M+H)

3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 11) $m/z = 802$ (M+H)

30 3-O-(4-Chlorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy)phenylmethyl]desosaminy-6-O-methyl erythronolide A (Compound no. 65) $m/z = 834$ (M+H)

3-O-(2-Thiophene)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy)phenylmethyl]desosaminy-6-O-methyl erythronolide A (Compound No. 62) $m/z = 806$ (M+H)

- 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 12) $m/z = 802$ (M+H)
- 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 13) $m/z = 802$ (M+H)
- 5 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 73) $m/z = 834$ (M+H)
- 3-O-(4-Methoxyphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 70) $m/z = 850$ (M+H)
- 10 3-O-(2,4-Difluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 74) $m/z = 820$ (M+H)
- 3-O-(2,4-Difluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 75) $m/z = 856$ (M+H)
- 15 3-O-(2-Bromophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 76) $m/z = 878$ (M+H), 880 (M+2+H)
- 3-O-(2-Bromophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 77) $m/z = 862$ (M+H), 864 (M+2+H)
- 20 3-O-(1-Naphthyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 71) $m/z = 850$ (M+H)
- 3-O-(1-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 72) $m/z = 834$ (M+H)
- 3-O-(2-Methylphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 66) $m/z = 814$ (M+H)
- 25 3-O-(2-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 67) $m/z = 798$ (M+H)
- 3-O-(4-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 68)
- 30 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 14) $m/z = 838$ (M+H)
- 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy)benzyl]desosaminy-6-O-methyl erythronolide A (Compound No. 17) $m/z = 818$ (M+H)
- 3-O-(4-Fluorophenyl)-5-O-(3'-N-desmethyl-3'-N-cyclopropyl) desosaminy-6-O-methyl erythronolide A (Compound No. 15) $m/z = 752$ (M+H)

3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 16) $m/z = 752$ (M+H)

3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl) desosaminyl-6-O-methyl erythronolide A (Compound No. 7) $m/z = 752$ (M+H)

5 Method B: 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-propargyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 18)

To a solution of 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl]desosaminyl-6-O-methyl erythronolide A (300 mg, 0.42 mmol, from step 5) in acetonitrile (10 ml) was added sodium bicarbonate (177 mg, 2.11 mmol) and propargyl bromide (69 mg, 0.46
10 mmol, 80% in toluene) and the mixture was stirred under nitrogen at 35°C-45°C for 24 to 30 hours. The reaction mixture was diluted with ethyl acetate and the solution was washed with 5% sodium bicarbonate solution, brine, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum. The product was purified by column chromatography using 8-10% acetone in hexane to obtain the desired product,
15 60mg (19%) $m/z = 750$ (M+H).

The following compounds were prepared similarly:

3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A. (Compound No. 22) $m/z = 750$ (M+H)

20 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A. (Compound No. 34) $m/z = 750$ (M+H)

EXAMPLE 5: Preparation of 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 21)

25 Step 3: 3-Hydroxy-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A

3-Hydroxy-5-O-(2'-O-benzoyl)desosaminyl-6-O-methyl erythronolide A (1.0 g, 1.44 mmol, example 1, step 2) was dissolved in dry acetonitrile (40 mL) and N-iodosuccinimide (422 mg, 1.87 mmol) was added at 0° C. The whole reaction mixture was stirred at ambient temperature for about 48 hours and then concentrated to dryness. To this
30 residue was added methanol (4 mL) and saturated ammonium chloride solution (4 mL). The whole reaction mixture was heated at 80° C for about 1 hour, pH of the solution was adjusted to 8. It was then extracted with ethyl acetate. Ethyl acetate layer was washed with

water, brine, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum. The crude product was then purified by column chromatography using hexane:acetone:triethylamine (6:1:0.1) to obtain the desired product, 540 mg (55%).

Step 4: 3-Hydroxy-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A

To a solution of 3-hydroxy-5-O-(2'-O-benzoyl-3'-N-desmethyl)desosaminyl-6-O-methyl erythronolide A (1.0 g, 1.47 mmol, from step 3) in acetonitrile (30 mL) was added sodium bicarbonate (NaHCO_3 , 515 mg, 6.13 mmol) and propargyl bromide (0.20 g, 1.68 mmol, 80% in toluene) and the mixture was stirred under nitrogen at 35 °C for about 14 hours. The reaction mixture was diluted with ethyl acetate and the solution was washed with 5% sodium bicarbonate (NaHCO_3) solution, brine, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum. The product was purified by column chromatography using 8–10% acetone in hexane to obtain the desired product, 650 mg (61.6%).

The following compounds were prepared analogously, using the appropriate bromide:

3-O-Hydroxy-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A

3-Hydroxy-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A

3-Hydroxy-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A

Step 5: 3-O-(2-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A

To a cold solution of 3-hydroxy-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (2 g, 2.78 mmol, from step 4) in dichloromethane were added DCC (1.4 g, 6.96 mmol), DMAP (0.85 g, 6.96 mmol) pyridine (1.12 g, 14.17 mmol) and 2-fluorophenyl acetic acid (1.1g, 6.96 mmol). The whole reaction mixture was stirred at 0 °C to 40 °C for about 16 hours. It was filtered through celite bed and washed with ethyl acetate. Organic layer was washed with saturated sodium bicarbonate (NaHCO_3) solution, water, brine, dried over anhydrous sodium sulphate (Na_2SO_4) and the solvent was removed under vacuum. The crude product was

then purified by column chromatography using 8–10% acetone in hexane to obtain the desired product, 1.8g (7.5%).

The following compounds were prepared analogously, using the appropriate acetic acid.

- 5 3-O-(3-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 10 3-O-(3-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Nitrophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(3-Nitrophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 15 3-O-(4-Pyridyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A

Step 6: 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Method A)

- 20 3-O-(2-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (300 mg, 0.36 mmol) was taken in methanol (10 mL) and heated with stirring for about 78 hours. Methanol was removed and the product was purified by column chromatography using 10 % acetone in hexane to obtain the desired compound, 90 mg (48%). $m/z = 733$ (M+H).

The following compounds were prepared similarly

- 25 3-O-(3-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 3-O-(4-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 30 3-O-(3-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A
- 35 3-O-(4-Nitrophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A

3-O-(3-Nitrophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A

5 3-O-(4-Pyridyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A

Step 6: 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Method A)

10 3-O-(2-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (300 mg, 0.36 mmol) was taken in methanol (10 mL) and heated with stirring for about 78 hours. Methanol was removed and the product was purified by column chromatography using 10 % acetone in hexane to obtain the desired compound, 90 mg (48%). $m/z = 733$ (M+H)

The following compounds were prepared similarly

15 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 20) $m/z = 752$ (M+H)

3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 35) $m/z = 777$ (M+H)

20 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-isopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 36) $m/z = 781$ (M+H)

3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 39) $m/z = 749$ (M+H)

3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 42) $m/z = 785$ (M+H)

25 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 43) $m/z = 785$ (M+H)

3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 44) $m/z = 735$ (M+H)

30 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 45) $m/z = 735$ (M+H)

3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 48) $m/z = 733$ (M+H)

3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 49) $m/z = 735$ (M+H)

3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 54) $m/z = 735$ (M+H)

Method B: 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 19)

5 3-O-(3-Fluorophenyl)acetyl-5-O-(2'-O-benzoyl-3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (1.6 g, 1.87 mmol) was taken in ethanol (130 ml) along with sodium bicarbonate (168 mg, 1.6 mmol) heated to 90°C to 95°C for 24 to 30 hours. Ethanol was removed and the product was purified by column chromatography to yield the desired product, 1.32 g (94.3%). $M/z = 752$ (M+H)

10 **Method C: 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-Desmethyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 23)**

3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-benzoyl-3'-N-allyl)desosaminyl-6-O-methyl erythronolide A (350 mg, 0.41 mmol) was taken in methanol (20 ml). To it was added sodium carbonate (86 mg, 0.82 mmol) and heated to 70°C to 75°C for 10 to 12
15 hours. Methanol was evaporated and the resulting compound was purified by column chromatography to yield the desired product, 100 mg (32.6%) (M+H)= 752.

The following compounds were prepared similarly:

3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 46) $m/z = 733$ (M+H)


20 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 47) $m/z = 733$ (M+H)

3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 37) $m/z = 779$ (M+H)

25 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 38) $m/z = 779$ (M+H)

3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 55) $m/z = 735$ (M+H)






















Table 1



















































Formula I




























(Formula I, wherein, $R^3=R''=CH_3$, $R'=H$, $U=V=OH$, and $Y=Z=O$)






















Compound No.	R^1	R^2
1		
2		
3		
4		
5		
6		
7		
















8		C_2H_5
9		C_2H_5
10		C_2H_5
11		
12		
13		
14		
15		
16		
17		
18		
19		

20		
21		
22		
23		
24		
25		
26		
27		
28		C_2H_5
29		C_2H_5
30		C_2H_5
31		

32		
33		
34		
35		
36		
37		
38		
39		
40		
41		C_2H_5
42		
43		
44		
45		

46		
47		
48		
49		
50		
51		C_2H_5
52		C_2H_5
53		
54		
55		
56		
57		
58		
59		
60		C_2H_5

61		
62		
63		C_2H_5
64		
65		
66		
67		
68		
69		
70		
71		

72		
73		
74		
75		
76		
77		
78		C_2H_5
79		C_2H_5
80		C_2H_5

***In vitro* Antibacterial activity**

Compounds of the invention displayed antibacterial activity when tested by the agar incorporation method (NCCLS agar dilution method). Minimum inhibitory concentrations ($\mu\text{g/ml}$) were obtained for representative compounds of the invention, which are given in Table 2.

Briefly, 1 mg/ml concentration of stock solutions of the compounds and erythromycin A, clarithromycin, telithromycin and clindamycin are prepared in dimethylsulfoxide. Drug dilutions and the agar plates are prepared by NCCLS method. Inoculum is prepared by direct colony suspensions in normal saline and adjusted to 0.5 McFarland turbidity and subsequently diluted as per NCCLS guidelines in order to obtain 10^4 CFU/spot. CFU/ml of a few randomly selected cultures is performed. The cultures are replicated on the agar plates using replicator (Denley's multipoint). The plates are incubated at $35 \pm 2^\circ\text{C}$ and in 5% CO_2 atmosphere (for respiratory and Haemophilus strains) for 18-24 hours at $35 \pm 2^\circ\text{C}$. Minimum inhibitory concentrations (MIC) are determined after 18-24 hours. Q. C. strains are also included in each run of the study. The cation content of Mueller Hinton agar is checked by performing disk dilution of 10 μg gentamicin disk against *Pseudomonas aeruginosa* ATCC 27853. The zone of inhibition should fall between 16-21 mm. The concentration of drug at which there is complete disappearance of growth spot or formation of less than 10 colonies per spot is considered as MIC.

While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention.

TABLE 2 -- Minimum Inhibitory Concentrations for Particular Compounds Disclosed Herein

C.. No.	S. <i>pneum</i> AB 14 (erm)	S. <i>pneum</i> AB 29 (erm)	S. <i>pneum</i> AB 30 (erm)	S. <i>pneum</i> 1256 (erm)	S. <i>pneum</i> 1275 (erm)	S. <i>pneum</i> AB 34 (mef)**	S. <i>pneum</i> CS 1687 (mef)	S. <i>pneum</i> 1251 (mef)	S. <i>pneum</i> 1294 (mef)	S. <i>pneum</i> 49619 ATCC	S. <i>pneum</i> 6303 ATCC	Mora- Xella MI	H.inflenza ATCC 49247
E	>16	>16	>16	>16	>16	>16	>16	8	>16	0.008	0.03	0.06	>16
C	>16	>16	>16	>16	>16	>16	>16	8	>16	0.008	0.008	0.125	8
T	0.25	0.25	0.25	0.25	0.008	0.125	0.125	0.5	0.5	0.008	0.015	0.03	2
Cl	>16				>16	0.03	0.06	0.03	0.03	0.015	0.06	2	
5	>8		>8	>8	>16	>8	2	2	2	2	2	>8	>128
16	>16		>16	16	>16	>16	4	8	1	4	1	>16	>16
17	>16	<0.5	>16	>16	>16	>16	1	1	1	<0.5	<0.5	8	>16
18	16	1	>16	>16	>16	1	1	1	1	1	1	16	>16
19	>16	2	>16	>16	>16	16	2	2	4	2	2	16	
20	>16		>16	16	>16	>16	1	1	1	0.25	0.5	16	>16
21	>16	>16	>16	>16	>16	>16	0.5	0.25	2	>16	0.5	0.25	>16
22	>16		>16	>16	>16	>16	1	4	4	0.5	0.5	4	>16
23	>16	1	>16	>16	>16	>16	2	2	2	2	2	16	>16
24	8	>16	>16	>16	>16	>16	0.5	1	2	0.5	0.5	2	
30	>16	<0.5	>16	>16	>16	>16	1	1	2	1	1	4	>16
50	>16	0.5	>16	>16	>16	>16	1	0.5	2	0.5	0.5	2	>16
73	>16	>16	>16	>16	>16	>16	2	2	>16	2	1	16	
74	16	>16	>16	>16	>16	>16	>16	>16	>16	>16	8	>16	

E: Erythromycin A, C: Clarithromycin, T: Telithromycin, Cl: Clindamycin, S. *pneum*: *streptococcus pneumoniae*

*


* Ribosomal modification (mutation) erm genes

**

** Efflux pump mef genes

WE CLAIM:

- 1 1. A compound having the structure of Formula I



7 **Formula I**

- 8 and its pharmaceutically acceptable acid addition salts, pharmaceutically acceptable
9 solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein

10 R^1 represents: lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) having one or more halogen
11 (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C_1-C_5) amino group, lower alkyl amino
12 (C_1-C_5) carbonyl group; lower alkoxy group (C_1-C_5); or five or six membered aryl or
13 heteroaryl ring having 1 to 3 hetero atoms selected from the group consisting of oxygen,
14 nitrogen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or
15 substituted by 1 to 3 substituents independently selected from the group consisting of
16 lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) group having one or more halogen (F, Cl,
17 Br, I) atoms, lower alkoxy (C_1-C_5) groups, lower alkyl (C_1-C_5) amino group, halogen
18 atoms (F, Cl, Br, I), amino group, nitro group, hydroxy group, and cyano group;

19 R^2 and R^3 are independently selected from: C_1-C_6 alkyl group optionally substituted with
20 halogen atoms (F, Cl, Br, I); cycloalkyl (C_3-C_7) group; or five to six membered aryl or
21 heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting
22 of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted
23 or substituted by 1 to 3 substituents independently selected from the group consisting of
24 lower alkyl (C_1-C_3), lower alkyl (C_1-C_3) group having one or more halogen (F, Cl, Br, I)
25 atom as substituent(s), lower alkoxy (C_1-C_3) group, lower alkyl (C_1-C_3) amino group,
26 halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, or cyano group; the above-
27 mentioned C_1-C_6 alkyl group may be substituted by: $NHCOR^5$, $NHCOOR^5$, $OCOR^5$,
28 COR^5 wherein R^5 represents lower alkyl (C_1-C_5); five to six membered aryl or heteroaryl
29 ring having 1 to 3 hetero atom independently selected from the group consisting of
30 nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or

31 substituted by 1 to 3 substituents independently selected from the group consisting of
32 lower alkyl (C₁-C₃), lower alkyl (C₁-C₃) group having one or more halogen (F, Cl, Br, I)
33 atoms as substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino group,
34 halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, and cyano group; C₂-C₆ alkenyl
35 or alkyne group optionally substituted with halogen (F, Cl, Br, I) atoms or a group
36 consisting of NHCOR⁵, NHCOOR⁵, COR⁵, OCOR⁵ (wherein R⁵ is as defined above);
37 cycloalkyl (C₃-C₇) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero
38 atom independently selected from the group consisting of nitrogen, oxygen, and sulphur,
39 wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3
40 substituents independently selected from the group consisting of lower alkyl (C₁-C₃)
41 group, lower alkyl (C₁-C₃) group having one or more halogen (F, Cl, Br, I) atoms as
42 substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino, halogen (F, Cl, Br,
43 I) atoms, nitro group, hydroxy group, amino group, and cyano group;

44 R' represents hydrogen, or a hydroxy protecting group optionally selected from acetyl,
45 benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxy methyl;

46 R'' represents hydrogen, or a lower alkyl (C₁-C₃) group;


47 Y represents oxygen or sulphur;

48 Z represents an oxygen atom or a group represented by NOR⁶, wherein R⁶ represents
49 hydrogen atom, alkyl (C₁-C₆) group, alkyl (C₁-C₆) amino group, phenyl or benzyl group,
50 or phenyl or benzyl group having 1 to 5 substituent independently selected from halogen
51 (F, Cl, Br, I) atoms, lower alkyl (C₁-C₃) group, hydroxy group, nitro group, cyano group,
52 or amino group;


53 U represents a hydroxy group: OR⁷, wherein R⁷ represents hydroxy protecting group
54 selected from acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxymethyl; or
55 -NH(CH₂)_nR⁸, wherein n represents 0 to 4 and R⁸ represents five or six membered aryl or
56 heteroaryl ring having 1 to 4 hetero atom independently selected from the group consisting
57 of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted
58 or substituted by one to three substituents independently selected from the group
59 consisting of lower alkyl (C₁-C₃) group, lower alkyl (C₁-C₃) group having one or more
60 halogen (F, Cl, Br, I) atoms as substituent (s), lower alkoxy (C₁-C₃) group, lower alkyl
61 (C₁-C₃) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, and
62 cyano group;

V represents: hydrogen atom; hydroxy group; or OR^7 , wherein R^7 represents a hydroxy protecting group selected from the group consisting of acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl and methoxymethyl;

U and V may also together represent (with carbon atoms at the 11- and 12- positions on the erythronolide skeleton): a group represented by Formula



or a group represented by the Formula



wherein R^9 represents: hydrogen atom; alkyl (C_1-C_6) group, wherein the alkyl (C_1-C_6) may be unsubstituted or substituted by halogen (F, Cl, Br, I) atoms, five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl (C_1-C_3) group, lower alkyl (C_1-C_3) group having one or more halogen (F, Cl, Br, I) atoms as substituent(s), lower alkoxy (C_1-C_3) group, lower alkyl (C_1-C_3) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group.

2. A compound selected from the group consisting of:

3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 1)

3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 2)

3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminyl erythronolide A (Compound No. 3)

3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluorobenzyl)desosaminyl-6-O-methyl erythronolide A (Compound No. 4)

- 11 3-O-(2-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-
12 6-O-methyl erythronolide A (Compound No. 5)
- 13 3-O-(3-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-fluoro)benzyl]desosaminyl-
14 6-O-methyl erythronolide A (Compound No. 6)
- 15 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl) desosaminyl-6-
16 O-methyl erythronolide A (Compound No. 7)
- 17 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl
18 erythronolide A (Compound No. 8)
- 19 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl
20 erythronolide A (Compound No. 9)
- 21 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminyl-6-O-methyl erythronolide A
22 (Compound No. 10)
- 23 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-
24 methyl erythronolide A (Compound No. 11)
- 25 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)
26 desosaminyl-6-O-methyl erythronolide A (Compound No. 12)
- 27 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)
28 desosaminyl-6-O-methyl erythronolide A (Compound No. 13)
- 29 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)benzyl]
30 desosaminyl-6-O-methyl erythronolide (Compound No. 14)
- 31 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-cyclopropyl] desosaminyl-6-
32 O-methyl erythronolide A (Compound No. 15)
- 33 3-O-(3-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-cyclopropyl]desosaminyl-6-O-
34 methyl erythronolide A (Compound No. 16)
- 35 3-O-(4-Fluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl]
36 desosaminyl-6-O-methyl erythronolide A (Compound No. 17)
- 37 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-
38 methyl erythronolide A (Compound No. 18)
- 39 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl
40 erythronolide A (Compound No. 19)
- 41 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminyl-6-O-methyl
42 erythronolide A (Compound No. 20)
- 43 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminyl-6-O-
44 methyl erythronolide A (Compound No. 21)

- 45 3-O-(3-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-
46 methyl erythronolide A (Compound No. 22)
- 47 3-O-(2-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl
48 erythronolide A (Compound No. 23)
- 49 3-O-(4-Fluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-(4-nitro)
50 benzyl)desosaminy-6-O-methyl erythronolide A (Compound No. 24)
- 51 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-
52 cyclopropylmethyl)desosaminy-6-O-methyl erythronolide A (Compound No. 25)
- 53 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)
54 desosaminy-6-O-methyl erythronolide A (Compound No. 26)
- 55 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminy-6-O-methyl
56 erythronolide A (Compound No. 27)
- 57 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl
58 erythronolide A (Compound No. 28)
- 59 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl
60 erythronolide A (Compound No. 29)
- 61 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl erythronolide A
62 (Compound No. 30)
- 63 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl
64 erythronolide A (Compound No. 31)
- 65 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl
66 erythronolide A (Compound No. 32)
- 67 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl
68 erythronolide A (Compound No. 33)
- 69 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-
70 methyl erythronolide A (Compound No. 34)
- 71 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-
72 methyl erythronolide A (Compound No. 35)
- 73 3-O-(2-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-isopropyl)desosaminy-6-O-
74 methyl erythronolide A (Compound No. 36)
- 75 3-O-(3-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-
76 methyl erythronolide A (Compound No. 37)
- 77 3-O-(4-Nitrophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-
78 methyl erythronolide A (Compound No. 38)


- 79 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminy-6-
80 O-methyl erythronolide A (Compound No. 39)
- 81 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminy-6-
82 O-methyl erythronolide A (Compound No. 40)
- 83 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl) desosaminy-6-O-methyl
84 erythronolide A (Compound No. 41)
- 85 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)
86 desosaminy-6-O-methyl erythronolide A (Compound No. 42)
- 87 3-O-(2-Pyridyl)acetyl-5-O-[3'-N-desmethyl-3'-N-benzyl]desosaminy-6-O-methyl
88 erythronolide A (Compound No. 43)
- 89 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl
90 erythronolide A (Compound No. 44)
- 91 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl
92 erythronolide A (Compound No. 45)
- 93 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl
94 erythronolide A (Compound No. 46)
- 95 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl
96 erythronolide A (Compound No. 47)
- 97 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-propargyl)desosaminy-6-O-methyl
98 erythronolide A (Compound No. 48)
- 99 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-
100 methyl erythronolide A (Compound No. 49)
- 101 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)desosaminy-6-
102 O-methyl erythronolide A (Compound No. 50)
- 103 3-O-(4-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl
104 erythronolide A (Compound No. 51)
- 105 3-O-(3-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desosaminy-6-O-methyl
106 erythronolide A (Compound No. 52)
- 107 3-O-(4-Pyridyl) acetyl-5-O-N-desmethyl-3'-N-cyclopropylmethyl) desosaminy-6-O-
108 methyl erythronolide A (Compound No. 53)
- 109 3-O-(4-Pyridyl) acetyl-5-O-(3'-N-desmethyl-3'-N-allyl)desosaminy-6-O-methyl
110 erythronolide A (Compound No. 54)
- 111 3-O-(2-Pyridyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropyl)desosaminy-6-O-
112 methyl erythronolide A (Compound No. 55)

- 113 3-O-(Phenyl)acetyl-5-O-[(3'-N-desmethyl-3'-N-cyclopropylmethyl]desoaminyl-6-O-
114 methyl erythronolide A (Compound No. 56)
- 115 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-(4-fluoro)benzyl)desoaminyl-6-O-
116 methyl erythronolide A (Compound No. 57)
- 117 3-O-(Phenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desoaminyl-6-O-methyl
118 erythronolide A (Compound No. 58)
- 119 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl) desosaminyl-
120 6-O-methyl erythronolide A (Compound No. 59)
- 121 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl
122 erythronolide A (Compound No. 60)
- 123 3-O-(2-Thiophene)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-methyl
124 erythronolide A (Compound No. 61)
- 125 3-O-(2-Thiophene)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl]
126 desosaminyl-6-O-methyl erythronolide A (Compound No. 62)
- 127 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl
128 erythronolide A (Compound No. 63)
- 129 3-O-(4-Chlorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)desosaminyl-6-O-
130 methyl erythronolide A (Compound No. 64)
- 131 3-O-(4-Chlorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3-hydroxy) benzyl]
132 desosaminyl-6-O-methyl erythronolide A (Compound No. 65)
- 133 3-O-(2-Methylphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]
134 desosaminyl-6-O-methyl erythronolide A (Compound No. 66)
- 135 3-O-(2-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)
136 desosaminyl-6-O-methyl erythronolide A (Compound No. 67)
- 137 3-O-(4-Methylphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)
138 desosaminyl-6-O-methyl erythronolide A (Compound No. 68)
- 139 3-O-(4-Methoxyphenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-cyclopropylmethyl)
140 desosaminyl-6-O-methyl erythronolide A (Compound No. 69)
- 141 3-O-(4-Methoxyphenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)
142 benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 70)
- 143 3-O-(1-Naphthyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]
144 desosaminyl-6-O-methyl erythronolide A (Compound No. 71)
- 145 3-O-(1-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)
146 desosaminyl-6-O-methyl erythronolide A (Compound No. 72)

- 147 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)
 148 desosaminyl-6-O-methyl erythronolide A (Compound No. 73)
- 149 3-O-(2,4-Difluorophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N-benzyl)
 150 desosaminyl-6-O-methyl erythronolide A (Compound No. 74)
- 151 3-O-(2,4-Difluorophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(3,4-difluoro)
 152 benzyl]desosaminyl-6-O-methyl erythronolide A (Compound No. 75)
- 153 3-O-(2-Bromophenyl)acetyl-5-O-[3'-N-desmethyl-3'-N-(4-hydroxy) benzyl]
 154 desosaminyl-6-O-methyl erythronolide A (Compound No. 76)
- 155 3-O-(2-Bromophenyl)acetyl-5-O-(3'-N-desmethyl-3'-N- benzyl)
 156 desosaminyl-6-O-methyl erythronolide A (Compound No. 77)
- 157 3-O-(3-Indole)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl
 158 erythronolide A (Compound No. 78)
- 159 3-O-(2-Naphthyl)acetyl-5-O-(3'-N-desmethyl-3'-N-ethyl)desoaminyl-6-O-methyl
 160 erythronolide A (Compound No. 79)

1 3. A pharmaceutical composition comprising a pharmaceutically effective amount of a
 2 compound as defined in claim 1 and 2 together with pharmaceutically acceptable
 3 carriers, excipients, or diluents.

1 4. A method for treating or preventing an animal or human suffering from bacterial
 2 infection caused by gram positive or gram negative or atypical pathogens comprising
 3 administering to a mammal in need of such treatment a pharmaceutically effective
 4 amount of a compound having the structure of Formula I,



14 **Formula I**

15 and its pharmaceutically acceptable acid addition salts, pharmaceutically acceptable
16 solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein

17 R^1 represents: lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) having one or more
18 halogen (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C_1-C_5) amino group, lower
19 alkyl amino (C_1-C_5) carbonyl group; lower alkoxy group (C_1-C_5); or five or six
20 membered aryl or heteroaryl ring having 1 to 3 hetero atoms selected from the group
21 consisting of oxygen, nitrogen, and sulphur, wherein the aryl or heteroaryl ring may be
22 unsubstituted or substituted by 1 to 3 substituents independently selected from the
23 group consisting of lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) group having one
24 or more halogen (F, Cl, Br, I) atoms, lower alkoxy (C_1-C_5) groups, lower alkyl (C_1-
25 C_5) amino group, halogen atoms (F, Cl, Br, I), amino group, nitro group, hydroxy
26 group, and cyano group;

27 R^2 and R^3 are independently selected from: C_1-C_6 alkyl group optionally substituted
28 with halogen atoms (F, Cl, Br, I); cycloalkyl (C_3-C_7) group; or five to six membered
29 aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the
30 group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring
31 may be unsubstituted or substituted by 1 to 3 substituents independently selected from
32 the group consisting of lower alkyl (C_1-C_3), lower alkyl (C_1-C_3) group having one or
33 more halogen (F, Cl, Br, I) atom as substituent(s), lower alkoxy (C_1-C_3) group, lower
34 alkyl (C_1-C_3) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, or
35 cyano group; the above-mentioned C_1-C_6 alkyl group may be substituted by:
36 $NHCOR^5$, $NHCOOR^5$, $OCOR^5$, COR^5 wherein R^5 represents lower alkyl (C_1-C_5); five
37 to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently
38 selected from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl
39 or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents
40 independently selected from the group consisting of lower alkyl (C_1-C_3), lower alkyl
41 (C_1-C_3) group having one or more halogen (F, Cl, Br, I) atoms as substituent(s), lower
42 alkoxy (C_1-C_3) group, lower alkyl (C_1-C_3) amino group, halogen (F, Cl, Br, I) atoms,
43 nitro group, hydroxy group, and cyano group; C_2-C_6 alkenyl or alkyne group
44 optionally substituted with halogen (F, Cl, Br, I) atoms or a group consisting of
45 $NHCOR^5$, $NHCOOR^5$, COR^5 , $OCOR^5$ (wherein R^5 is as defined above); cycloalkyl
46 (C_3-C_7) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom

47 independently selected from the group consisting of nitrogen, oxygen, and sulphur,
48 wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3
49 substituents independently selected from the group consisting of lower alkyl (C₁-C₃)
50 group, lower alkyl (C₁-C₃) group having one or more halogen (F, Cl, Br, I) atoms as
51 substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino, halogen (F, Cl,
52 Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group;

53 R' represents hydrogen, or a hydroxy protecting group optionally selected from acetyl,
54 benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxy methyl;

55 R'' represents hydrogen, or a lower alkyl (C₁-C₃) group;


56 Y represents oxygen or sulphur;

57 Z represents an oxygen atom or a group represented by NOR⁶, wherein R⁶ represents
58 hydrogen atom, alkyl (C₁-C₆) group, alkyl (C₁-C₆) amino group, phenyl or benzyl
59 group, or phenyl or benzyl group having 1 to 5 substituent independently selected from
60 halogen (F, Cl, Br, I) atoms, lower alkyl (C₁-C₃) group, hydroxy group, nitro group,
61 cyano group, or amino group;


62 U represents a hydroxy group: OR⁷, wherein R⁷ represents hydroxy protecting group
63 selected from acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or
64 methoxymethyl; or -NH(CH₂)_nR⁸, wherein n represents 0 to 4 and R⁸ represents five or
65 six membered aryl or heteroaryl ring having 1 to 4 hetero atom independently selected
66 from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or
67 heteroaryl ring may be unsubstituted or substituted by one to three substituents
68 independently selected from the group consisting of lower alkyl (C₁-C₃) group, lower
69 alkyl (C₁-C₃) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s),
70 lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino, halogen (F, Cl, Br, I) atoms,
71 nitro group, hydroxy group, amino group, and cyano group;

72 V represents: hydrogen atom; hydroxy group; or OR⁷, wherein R⁷ represents a
73 hydroxy protecting group selected from the group consisting of acetyl, benzoyl,
74 butyldiphenylsilyl, methylthiomethyl and methoxymethyl;

U and V may also together represent (with carbon atoms at the 11- and 12- positions on the erythronolide skeleton): a group represented by Formula



or a group represented by the Formula



wherein R^9 represents: hydrogen atom; alkyl (C_1 - C_6) group, wherein the alkyl (C_1 - C_6) may be unsubstituted or substituted by halogen (F, Cl, Br, I) atoms, five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl (C_1 - C_3) group, lower alkyl (C_1 - C_3) group having one or more halogen (F, Cl, Br, I) atoms as substituent(s), lower alkoxy (C_1 - C_3) group, lower alkyl (C_1 - C_3) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group.


5. A method for treating or preventing of animal or human suffering from bacterial infections according to claim 4 caused by bacteria selected from the group consisting of *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Haemophilus influenzae*.

6. A method for treating or preventing an animal or human suffering from bacterial infection caused by gram positive or gram negative or atypical pathogens comprising administering to a mammal in need of such treatment therapeutically effective amount of a pharmaceutical composition according to claim 3.

7. A method for treating or preventing of animal or human suffering from bacterial infections caused by bacteria selected from the group consisting of *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Haemophilus influenzae*, comprising administering to a

mammal in need of such treatment therapeutically amount of a pharmaceutical composition according to claim 3.

8. A process for preparing a compound of Formula I



Formula I

and its pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein

$R^3 = R'' = CH_3$, $R' = H$, $U = V = OH$, and $Y = Z = O$

R^1 represents: lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C_1-C_5) amino group, lower alkyl amino (C_1-C_5) carbonyl group; lower alkoxy group (C_1-C_5); or five or six membered aryl or heteroaryl ring having 1 to 3 hetero atoms selected from the group consisting of oxygen, nitrogen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) group having one or more halogen (F, Cl, Br, I) atoms, lower alkoxy (C_1-C_5) groups, lower alkyl (C_1-C_5) amino group, halogen atoms (F, Cl, Br, I), amino group, nitro group, hydroxy group, and cyano group;

R^2 and R^3 are independently selected from: C_1-C_6 alkyl group optionally substituted with halogen atoms (F, Cl, Br, I); cycloalkyl (C_3-C_7) group; or five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from

29 the group consisting of lower alkyl (C₁-C₃), lower alkyl (C₁-C₃) group having one or
30 more halogen (F, Cl, Br, I) atom as substituent(s), lower alkoxy (C₁-C₃) group, lower
31 alkyl (C₁-C₃) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, or
32 cyano group; the above-mentioned C₁-C₆ alkyl group may be substituted by:
33 NHCOR⁵, NHCOOR⁵, OCOR⁵, COR⁵ wherein R⁵ represents lower alkyl (C₁-C₃); five
34 to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently
35 selected from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl
36 or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents
37 independently selected from the group consisting of lower alkyl (C₁-C₃), lower alkyl
38 (C₁-C₃) group having one or more halogen (F, Cl, Br, I) atoms as substituent(s), lower
39 alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino group, halogen (F, Cl, Br, I) atoms,
40 nitro group, hydroxy group, and cyano group; C₂-C₆ alkenyl or alkyne group
41 optionally substituted with halogen (F, Cl, Br, I) atoms or a group consisting of
42 NHCOR⁵, NHCOOR⁵, COR⁵, OCOR⁵ (wherein R⁵ is as defined above); cycloalkyl
43 (C₃-C₇) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom
44 independently selected from the group consisting of nitrogen, oxygen, and sulphur,
45 wherein the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3
46 substituents independently selected from the group consisting of lower alkyl (C₁-C₃)
47 group, lower alkyl (C₁-C₃) group having one or more halogen (F, Cl, Br, I) atoms as
48 substituent(s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino, halogen (F, Cl,
49 Br, I) atoms, nitro group, hydroxy group, amino group, and cyano group;

50 R' represents hydrogen, or a hydroxy protecting group optionally selected from acetyl,
51 benzoyl, butyldiphenylsilyl, methylthiomethyl, or methoxy methyl;

52 R'' represents hydrogen, or a lower alkyl (C₁-C₃) group;


53 Y represents oxygen or sulphur;

54 Z represents an oxygen atom or a group represented by NOR⁶, wherein R⁶ represents
55 hydrogen atom, alkyl (C₁-C₆) group, alkyl (C₁-C₆) amino group, phenyl or benzyl
56 group, or phenyl or benzyl group having 1 to 5 substituent independently selected from
57 halogen (F, Cl, Br, I) atoms, lower alkyl (C₁-C₃) group, hydroxy group, nitro group,
58 cyano group, or amino group;


59 U represents a hydroxy group: OR^7 , wherein R^7 represents hydroxy protecting group
 60 selected from acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, or
 61 methoxymethyl; or $-NH(CH_2)_nR^8$, wherein n represents 0 to 4 and R^8 represents five or
 62 six membered aryl or heteroaryl ring having 1 to 4 hetero atom independently selected
 63 from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or
 64 heteroaryl ring may be unsubstituted or substituted by one to three substituents
 65 independently selected from the group consisting of lower alkyl (C_1-C_3) group, lower
 66 alkyl (C_1-C_3) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s),
 67 lower alkoxy (C_1-C_3) group, lower alkyl (C_1-C_3) amino, halogen (F, Cl, Br, I) atoms,
 68 nitro group, hydroxy group, amino group, and cyano group;

69 V represents: hydrogen atom; hydroxy group; or OR^7 , wherein R^7 represents a
 70 hydroxy protecting group selected from the group consisting of acetyl, benzoyl,
 71 butyldiphenylsilyl, methylthiomethyl and methoxymethyl;

72 U and V may also together represent (with carbon atoms at the 11- and 12- positions
 73 on the erythronolide skeleton): a group represented by Formula




76 or a group represented by the Formula




81 wherein R^9 represents: hydrogen atom; alkyl (C_1-C_6) group, wherein the alkyl (C_1-C_6)
 82 may be unsubstituted or substituted by halogen (F, Cl, Br, I) atoms, five or six
 83 membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected
 84 from the group consisting of nitrogen, oxygen, and sulphur, wherein the aryl or
 85 heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents
 86 independently selected from the group consisting of lower alkyl (C_1-C_3) group, lower
 87 alkyl (C_1-C_3) group having one or more halogen (F, Cl, Br, I) atoms as substituent(s),
 88 lower alkoxy (C_1-C_3) group, lower alkyl (C_1-C_3) amino, halogen (F, Cl, Br, I) atoms,
 89 nitro group, hydroxy group, amino group, and cyano group, which method comprises:

Step (1) treating clarithromycin of Formula II




Formula II

with an acid at ambient temperature to give a compound of Formula III,




Formula III

Step (2) reacting the compound of Formula III with a reagent of Formula R'_2O or $R'X$ (wherein R' is hydroxy protecting group optionally selected from acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, methoxy methyl and X is an optional halogen atom) to give a compound of Formula IV,




Formula IV

Step (3) reacting the compound of Formula IV with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 (wherein R^1 is as defined for Formula I in claim 1 and R^4 is a group selected from pivaloyl group, p-toluenesulfonyl group, isobutoxycarbonyl group, ethoxycarbonyl group or isopropoxycarbonyl group) to give a compound of Formula V,




Formula V

Step (4) treating the compound of Formula V with aqueous alcohol to give a compound of Formula VI,



Formula VI


Step (5) desmethylating at 3'-N-dimethyl group of the compound of Formula VI with N-iodosuccinamide and acetonitrile or iodine in presence of sodium acetate followed by quench with sodium thiosulphate to give a compound of Formula VII,



Formula VII

- 1 13. The process according to claim 8 wherein, the reaction of compound of Formula IV
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula V is carried out in presence of an inorganic base selected from
4 the group comprising of sodium hydrogen carbonate, potassium carbonate or organic
5 base selected from the group comprising of triethylamine, pyridine, tributylamine and
6 4-dimethylaminopyridine.
- 1 14. The process according to claim 8 wherein, the reaction of compound of Formula IV
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula V is carried out in presence of an inert solvent selected from the
4 group comprising of dichloromethane, dichloroethane, acetone, ethyl acetate and
5 tetrahydrofuran.
- 1 15. The process according to claim 8 wherein, the reaction of compound of Formula V is
2 carried out with aqueous alcohol selected from the group comprising of aqueous
3 methanol, aqueous ethanol, aqueous propanol and aqueous isopropanol to give a
4 compound of Formula VI.
- 1 16. The process according to claim 8 wherein, the reaction of the compound of Formula
2 VII with a reagent of Formula R^2CHO or R^2_2CO to give a compound of Formula I is
3 carried out in presence of a reducing agent selected from the group comprising of
4 sodium borohydride, sodium cyanoborohydride, sodium triacetoborohydride or
5 palladium/carbon catalyst.
- 1 17. The process according to claim 8 wherein, the reaction of the compound of Formula
2 VII with a reagent of Formula R^2CHO or R^2_2CO to give a compound of Formula I is
3 carried out in presence of a protic or non-protic solvent selected from the group
4 comprising of hexane, toluene, methylene chloride, ethylene chloride, chloroform,
5 tetrahydrofuran, N-methyl-pyrrolidinone, diethyl ether, bis-methoxymethyl ether,
6 dimethylformamide, acetonitrile, acetone and ethyl acetate.

18. A Process for preparing a compound of Formula I



Formula I

and its pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein


$R^3=R''=CH_3$, $R'=H$, $U=V=OH$, $Y=Z=O$

R^1 represents lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C_1-C_5) amino group, lower alkyl amino (C_1-C_5) carbonyl group, lower alkoxy group (C_1-C_5), five or six membered aryl or heteroaryl ring having 1 to 3 hetero atoms selected from the group consisting of oxygen, nitrogen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) group having one or more halogen (F, Cl, Br, I) atoms, lower alkoxy (C_1-C_5) groups, lower alkyl (C_1-C_5) amino group, halogen atoms (F, Cl, Br, I), amino group, nitro group, hydroxy group, cyano group;


R^2 is selected from C_1-C_6 alkyl group optionally substituted with halogen atoms (F, Cl, Br, I), cycloalkyl (C_3-C_7) group, five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl (C_1-C_3), lower alkyl (C_1-C_3) group having one or more halogen (F, Cl, Br, I) atom as substituent (s), lower alkoxy (C_1-C_3) group, lower alkyl (C_1-C_3) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, cyano group, C_1-C_6 alkyl group may also be substituted by a group consisting of $NHCOR^5$, $NHCOOR^5$, $OCOR^5$, COR^5 [wherein R^5 represents lower alkyl (C_1-C_5), five to six membered aryl or heteroaryl

ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl (C₁-C₃), lower alkyl (C₁-C₃) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, cyano group]; C₂-C₆ alkenyl or alkyne group optionally substituted with halogen (F, Cl, Br, I) atoms or a group consisting of NHCOR⁵, NHCOOR⁵, COR⁵, OCOR⁵ (wherein R⁵ is as defined above); cycloalkyl (C₃-C₇) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl (C₁-C₃) group, lower alkyl (C₁-C₃) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkoxy (C₁-C₃) group, lower alkyl (C₁-C₃) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, cyano group, which method comprises the steps of


Step (1) desmethylating at 3'-N-dimethyl group of the compound of Formula II with N-iodosuccinamide and acetonitrile or iodine in presence of sodium acetate followed by quench with sodium thiosulphate to give a compound of Formula VIII




Formula II

**Formula VIII**

Step (2) reacting the compound of Formula VIII with a reagent of Formula R^2CHO or R^2CO (wherein R^2 is as defined for Formula 1 in claim 1) to give a compound of Formula IX


**Formula IX**

Step (3) treating the compound of Formula IX with acid at an ambient temperature to give a compound of Formula X

**Formula X**


Step (4) reacting the compound of Formula X with a reagent of Formula $R'O$ or $R'X$ (wherein R' is hydroxy protecting group optionally selected from acetyl, benzoyl,

butyldiphenylsilyl, methylthiomethyl, methoxy methyl and X is an optional halogen atom) to give a compound of Formula XI




Formula XI

Step (5) reacting the compound of Formula XI with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 (wherein R^1 is as defined for Formula I in claim 1 and R^4 is a group selected from pivaloyl group, p-toluenesulfonyl group, isobutoxycarbonyl group, ethoxycarbonyl group or isopropoxycarbonyl group) to give a compound of Formula XII



Formula XII

Step (6) treating the compound of Formula XII with aqueous alcohol to give a compound of Formula I



Formula I

$R^3=R''=CH_3$, $R'=H$, $U=V=OH$, and $Y=Z=O$

19. The process according to claim 18 wherein, the reaction of the compound of Formula VIII with a reagent of Formula R^2CHO or R^2_2CO to give a compound of Formula IX is carried out in presence of a reducing agent selected from the group comprising of sodium borohydride, sodium cyanoborohydride, sodium triacetoborohydride or palladium/carbon catalyst.


20. The process according to claim 18 wherein, the reaction of the compound of Formula VIII with a reagent of Formula R^2CHO or R^2_2CO to give a compound of Formula IX is carried out in presence of a protic or non-protic solvent selected from the group comprising of hexane, toluene, methylene chloride, ethylene chloride, chloroform, tetrahydrofuran, N-methyl-pyrrolidinone, diethyl ether, bis-methoxymethyl ether, dimethylformamide, acetonitrile, acetone and ethyl acetate.

21. The process according to claim 18 wherein, the reaction of compound of Formula IX with hydrochloric or dichloroacetic acid is carried out with aqueous alcohol selected from the group comprising of aqueous methanol, aqueous ethanol, aqueous propanol and aqueous isopropanol to give a compound of Formula X.

22. The process according to claim 18 wherein, the reaction of compound of Formula X with a reagent of Formula R'_2O or $R'X$ to give a compound of Formula XI is carried out in presence of an inorganic base selected from the group comprising of sodium hydrogen carbonate, potassium carbonate or an organic base selected from the group comprising of triethylamine, pyridine, tributylamine and 4-N-dimethylamonipyridine.

- 1 23. The process according to claim 18 wherein, the reaction of compound of Formula X
2 with a reagent of Formula R'_2O or $R'X$ to give a compound of Formula XI is carried
3 out in presence of an inert solvent selected from the group comprising of
4 dichloromethane, dichloroethane, acetone, ethyl acetate and tetrahydrofuran.
- 1 24. The process according to claim 18 wherein, the reaction of compound of Formula XI
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula XII is carried out in presence of an activating agent selected
4 from the group comprising of dichlorohexylcarbodiimide (DCC) and 1-ethyl-3(3-
5 dimethylaminopropyl) carbodiimide hydrochloride (EDCI).
- 1 25. The process according to claim 18 wherein, the reaction of compound of Formula XI
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula XII is carried out in presence of an inorganic base selected from
4 the group comprising of sodium hydrogen carbonate, potassium carbonate or organic
5 base selected from the group comprising of triethylamine, pyridine, tributylamine and
6 4-dimethylaminopyridine.
- 1 26. The process according to claim 18 wherein, the reaction of compound of Formula XI
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula XII is carried out in presence of an inert solvent selected from
4 the group comprising of dichloromethane, dichloroethane, acetone, ethyl acetate and
5 tetrahydrofuran.
- 1 27. The process according to claim 18 wherein, the reaction of compound of Formula XII
2 is carried out with aqueous alcohol selected from the group comprising of aqueous
3 methanol, aqueous ethanol, aqueous propanol and aqueous isopropanol to give a
4 compound of Formula I.

28. A process for preparing a compound of Formula I



Formula I

and its pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein


$R^3=R''=CH_3$, $R'=H$, $U=V=OH$, $Y=Z=O$

R^1 represents lower alkyl (C_1 - C_5) group, lower alkyl (C_1 - C_5) having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C_1 - C_5) amino group, lower alkyl amino (C_1 - C_5) carbonyl group, lower alkoxy group (C_1 - C_5), five or six membered aryl or heteroaryl ring having 1 to 3 hetero atoms selected from the group consisting of oxygen, nitrogen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl (C_1 - C_5) group, lower alkyl (C_1 - C_5) group having one or more halogen (F, Cl, Br, I) atoms, lower alkoxy (C_1 - C_5) groups, lower alkyl (C_1 - C_5) amino group, halogen atoms (F, Cl, Br, I), amino group, nitro group, hydroxy group, cyano group;

R^2 is selected from C_1 - C_6 alkyl group optionally substituted with halogen atoms (F, Cl, Br, I), cycloalkyl (C_3 - C_7) group, five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl (C_1 - C_3), lower alkyl (C_1 - C_3) group having one or more halogen (F, Cl, Br, I) atom as substituent (s), lower alkoxy (C_1 - C_3) group, lower alkyl (C_1 - C_3) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, cyano group, C_1 - C_6 alkyl group may


also be substituted by a group consisting of NHCOR^5 , NHCOOR^5 , OCOR^5 , COR^5 [wherein R^5 represents lower alkyl ($\text{C}_1\text{-C}_3$), five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl ($\text{C}_1\text{-C}_3$), lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, cyano group]; $\text{C}_2\text{-C}_6$ alkenyl or alkyne group optionally substituted with halogen (F, Cl, Br, I) atoms or a group consisting of NHCOR^5 , NHCOOR^5 , COR^5 , OCOR^5 (wherein R^5 is as defined above); cycloalkyl ($\text{C}_3\text{-C}_7$) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, cyano group, which method comprises the steps of

Step (1) treating clarithromycin of Formula II




Formula II

with acid at ambient temperature to give a compound of Formula III



Formula III


Step (2) reacting the compound of Formula III with a reagent of Formula R'_2O or $R'X$ (wherein R' is hydroxy protecting group optionally selected from acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, methoxy methyl and X is an optional halogen atom) to give a compound of Formula IV



Formula IV


Step (3) reacting the compound of Formula IV with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 (wherein R^1 is as defined for Formula I in claim 1 and R^4 is a group selected from pivaloyl group, p-toluenesulfonyl group, isobutoxycarbonyl group, ethoxycarbonyl group or isopropoxycarbonyl group) to give a compound of Formula V

78




Formula V

Step (4) desmethylating at 3'-N-dimethyl group of the compound of Formula V with N-iodosuccinamide and acetonitrile or iodine in presence of sodium acetate followed by quench with sodium thiosulphate to obtain the compound of Formula XIII




Formula XIII

Step (5) treating the compound of Formula XIII with aqueous alcohol to give a compound of Formula VII



Formula VII

Step (6) reacting the compound of Formula VII with a reagent of Formula $R^2\text{CHO}$ or $R^2_2\text{CO}$ (wherein R^2 is as defined for Formula 1 in claim 1) to give a compound of Formula I



Formula I

$R^3=R''=\text{CH}_3$, $R'=\text{H}$, $\text{U}=\text{V}=\text{OH}$, and $\text{Y}=\text{Z}=\text{O}$

29. The process according to claim 28 wherein, the reaction of clarithromycin of Formula II with hydrochloric or dichloroacetic acid to give a compound of Formula III is carried out in presence of aqueous alcohol selected from the group comprising of aqueous methanol, aqueous ethanol, aqueous propanol and aqueous isopropanol.


30. The process according to claim 28 wherein, the reaction of compound of Formula III with a reagent of Formula $R'_2\text{O}$ or $R'X$ to give a compound of Formula IV is carried out in presence of an inorganic base selected from the group comprising of sodium hydrogen carbonate, potassium carbonate or an organic base selected from the group comprising of triethylamine, pyridine, tributylamine and 4-N-dimethylamonipyridine.

31. The process according to claim 28 wherein, the reaction of compound of Formula III with a reagent of Formula $R'_2\text{O}$ or $R'X$ to give a compound of Formula IV is carried out in presence of an inert solvent selected from the group comprising of dichloromethane, dichloroethane, acetone, ethyl acetate and tetrahydrofuran.

32. The process according to claim 28 wherein, the reaction of compound of Formula IV with a reagent of Formula $R^1\text{COOH}$, $R^1\text{COX}$, $(R^1\text{CO})_2\text{O}$ or $R^1\text{COOR}^4$ to give a compound of Formula V is carried out in presence of an activating agent selected from the group comprising of dichlorohexylcarbodiimide (DCC) and 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDCI).

- 1 33. The process according to claim 28 wherein, the reaction of compound of Formula IV
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula V is carried out in presence of an inorganic base selected from
4 the group comprising of sodium hydrogen carbonate, potassium carbonate or organic
5 base selected from the group comprising of triethylamine, pyridine, tributylamine and
6 4-dimethylaminopyridine.
- 1 34. The process according to claim 28 wherein, the reaction of compound of Formula IV
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula V is carried out in presence of an inert solvent selected from the
4 group comprising of dichloromethane, dichloroethane, acetone, ethyl acetate and
5 tetrahydrofuran.
- 1 35. The process according to claim 28 wherein, the reaction of compound of Formula XIII
2 is carried out with aqueous alcohol selected from the group comprising of aqueous
3 methanol, aqueous ethanol, aqueous propanol and aqueous isopropanol to give a
4 compound of Formula VII.
- 1 36. The process according to claim 28 wherein, the reaction of the compound of Formula
2 VII with a reagent of Formula R^2CHO or R^2_2CO to give a compound of Formula I is
3 carried out in presence of a reducing agent selected from the group comprising of
4 sodium borohydride, sodium cyanoborohydride, sodium triacetoborohydride or
5 palladium/carbon catalyst.
- 1 37. The process according to claim 28 wherein, the reaction of the compound of Formula
2 VII with a reagent of Formula R^2CHO or R^2_2CO to give a compound of Formula I is
3 carried out in presence of a protic or non-protic solvent selected from the group
4 comprising of hexane, toluene, methylene chloride, ethylene chloride, chloroform,
5 tetrahydrofuran, N-methyl-pyrrolidinone, diethyl ether, bis-methoxymethyl ether,
6 dimethylformamide, acetonitrile, acetone and ethyl acetate.

38. A process for preparing a compound of Formula I



Formula I

and its pharmaceutically acceptable acid addition salts, pharmaceutically acceptable solvates, enantiomers, diastereomers, polymorphs and metabolites, wherein


$R^3=R''=CH_3$, $R'=H$, $U=V=OH$, and $Y=Z=O$

R^1 represents lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkyl (C_1-C_5) amino group, lower alkyl amino (C_1-C_5) carbonyl group, lower alkoxy group (C_1-C_5), five or six membered aryl or heteroaryl ring having 1 to 3 hetero atoms selected from the group consisting of oxygen, nitrogen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl (C_1-C_5) group, lower alkyl (C_1-C_5) group having one or more halogen (F, Cl, Br, I) atoms, lower alkoxy (C_1-C_5) groups, lower alkyl (C_1-C_5) amino group, halogen atoms (F, Cl, Br, I), amino group, nitro group, hydroxy group, cyano group;

R^2 is selected from C_1-C_6 alkyl group optionally substituted with halogen atoms (F, Cl, Br, I), cycloalkyl (C_3-C_7) group, five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl (C_1-C_3), lower alkyl (C_1-C_3) group having one or more halogen (F, Cl, Br, I) atom as substituent (s), lower alkoxy (C_1-C_3) group, lower alkyl (C_1-C_3) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, cyano group, C_1-C_6 alkyl group may


also be substituted by a group consisting of NHCOR^5 , NHCOOR^5 , OCOR^5 , COR^5 [wherein R^5 represents lower alkyl ($\text{C}_1\text{-C}_5$), five to six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituent independently selected from the group consisting of lower alkyl ($\text{C}_1\text{-C}_3$), lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino group, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, cyano group]; $\text{C}_2\text{-C}_6$ alkenyl or alkyne group optionally substituted with halogen (F, Cl, Br, I) atoms or a group consisting of NHCOR^5 , NHCOOR^5 , COR^5 , OCOR^5 (wherein R^5 is as defined above); cycloalkyl ($\text{C}_3\text{-C}_7$) group; five or six membered aryl or heteroaryl ring having 1 to 3 hetero atom independently selected from the group consisting of nitrogen, oxygen, sulphur, the aryl or heteroaryl ring may be unsubstituted or substituted by 1 to 3 substituents independently selected from the group consisting of lower alkyl ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) group having one or more halogen (F, Cl, Br, I) atoms as substituent (s), lower alkoxy ($\text{C}_1\text{-C}_3$) group, lower alkyl ($\text{C}_1\text{-C}_3$) amino, halogen (F, Cl, Br, I) atoms, nitro group, hydroxy group, amino group, cyano group, which method comprises the steps of

Step (1) treating clarithromycin of Formula II




Formula II

with acid at ambient temperature to give a compound of Formula III




Formula III

Step (2) reacting the compound of Formula III with a reagent of Formula R'_2O or $R'X$ (wherein R' is hydroxy protecting group optionally selected from acetyl, benzoyl, butyldiphenylsilyl, methylthiomethyl, methoxy methyl and X is an optional halogen atom) to give a compound of Formula IV




Formula IV

Step (3) desmethylating at 3'-N-dimethyl group of the compound of Formula IV with N-iodosuccinamide and acetonitrile or iodine in presence of sodium acetate followed by quench with sodium thiosulphate to give a compound of Formula XIV




Formula XIV

Step (4) reacting the compound of Formula XIV with a reagent of Formula R^2CHO or R^2CO (wherein R^2 is as defined for Formula 1) to give a compound of Formula XI




Formula XI

Step (5) reacting the compound of Formula XI with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 (wherein R^1 is as defined for Formula I and R^4 is a group selected from pivaloyl group, p-toluenesulfonyl group, isobutoxycarbonyl group, ethoxycarbonyl group or isopropoxycarbonyl group) to give a compound of Formula XII



Formula XII

Step (6) treating the compound of Formula XII with aqueous alcohol to give the compound of Formula I



Formula I

$R^3=R''=CH_3$, $R'=H$, $U=V=OH$, and $Y=Z=O$

- 1 39. The process according to claim 38 wherein, the reaction of clarithromycin of Formula
- 2 II with hydrochloric or dichloroacetic acid to give a compound of Formula III is
- 3 carried out in presence of aqueous alcohol selected from the group comprising of
- 4 aqueous methanol, aqueous ethanol, aqueous propanol and aqueous isopropanol.

- 1 40. The process according to claim 38 wherein, the reaction of compound of Formula III
- 2 with a reagent of Formula R'_2O or $R'X$ to give a compound of Formula IV is carried
- 3 out in presence of an inorganic base selected from the group comprising of sodium
- 4 hydrogen carbonate, potassium carbonate or an organic base selected from the group
- 5 comprising of triethylamine, pyridine, tributylamine and 4-N-dimethylaminopyridine.

- 1 41. The process according to claim 38 wherein, the reaction of compound of Formula III
2 with a reagent of Formula R'_2O or $R'X$ to give a compound of Formula IV is carried
3 out in presence of an inert solvent selected from the group comprising of
4 dichloromethane, dichloroethane, acetone, ethyl acetate and tetrahydrofuran.
- 1 42. The process according to claim 38 wherein, the reaction of the compound of Formula
2 XIV with a reagent of Formula R^2CHO or R^2CO to give a compound of Formula XI
3 is carried out in presence of a reducing agent selected from the group comprising of
4 sodium borohydride, sodium cyanoborohydride, sodium triacetoborohydride or
5 palladium/carbon catalyst.
- 1 43. The process according to claim 38 wherein, the reaction of the compound of Formula
2 XIV with a reagent of Formula R^2CHO or R^2CO to give a compound of Formula XI
3 is carried out in presence of a protic or non-protic solvent selected from the group
4 comprising of hexane, toluene, methylene chloride, ethylene chloride, chloroform,
5 tetrahydrofuran, N-methyl-pyrrolidinone, diethyl ether, bis-methoxymethyl ether,
6 dimethylformamide, acetonitrile, acetone and ethyl acetate.
- 1 44. The process according to claim 38 wherein, the reaction of compound of Formula XI
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula XII is carried out in presence of an activating agent selected
4 from the group comprising of dichlorohexylcarbodiimide (DCC) and 1-ethyl-3(3-
5 dimethylaminopropyl) carbodiimide hydrochloride (EDCI).
- 1 45. The process according to claim 38 wherein, the reaction of compound of Formula XI
2 with a reagent of Formula R^1COOH , R^1COX , $(R^1CO)_2O$ or R^1COOR^4 to give a
3 compound of Formula XII is carried out in presence of an inorganic base selected from
4 the group comprising of sodium hydrogen carbonate, potassium carbonate or organic
5 base selected from the group comprising of triethylamine, pyridine, tributylamine and
6 4-dimethylaminopyridine.

- 1 46. The process according to claim 38 wherein, the reaction of compound of Formula XI
2 with a reagent of Formula $R^1\text{COOH}$, $R^1\text{COX}$, $(R^1\text{CO})_2\text{O}$ or $R^1\text{COOR}^4$ to give a
3 compound of Formula XII is carried out in presence of an inert solvent selected from
4 the group comprising of dichloromethane, dichloroethane, acetone, ethyl acetate and
5 tetrahydrofuran.
- 1 47. The process according to claim 28 wherein, the reaction of compound of Formula XII
2 is carried out with aqueous alcohol selected from the group comprising of aqueous
3 methanol, aqueous ethanol, aqueous propanol and aqueous isopropanol to give a
4 compound of Formula I.

INTERNATIONAL SEARCH REPORT

PCT/IB 03/04166

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07H17/08 A61K31/7048		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07H A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, CHEM ABS Data, PAJ, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 923 784 A (KIERSTEAD RICHARD W ET AL) 2 December 1975 (1975-12-02) the whole document	1-47
X	LEMAHIEU R A: "GLYCOSIDE CLEAVAGE REACTIONS OF ERYTHROMYCIN A. PREPARATION OF ERYTHRONOLIDE A" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 17, no. 9, 1 September 1974 (1974-09-01), pages 953-956, XP000651522 ISSN: 0022-2623 page compounds 5-6	1-47
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "G" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
23 March 2004		06/04/2004
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3018		Authorized officer Klein, D

INTERNATIONAL SEARCH REPORT

PCT/IB 03/04166

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TANIKAWA T. ET AL: "Synthesis and antibacterial activity of acylides (3-O-Acyl-erythromycin derivatives): A novel class of macrolide antibiotics" JOURNAL OF MEDICINAL CHEMISTRY, vol. 44, no. 24, 2001, pages 4027-4030, XP002271659 page 4028 compounds 2,3a,3c,3k	1-47
X	TANIKAWA, TETSUYA ET AL: "Synthesis and Antibacterial Activity of a Novel Series of Acylides: 3-O-(3-Pyridyl)acetylerythromycin A Derivatives" JOURNAL OF MEDICINAL CHEMISTRY (2003), 46(13), 2706-2715 , 2003, XP002271660 page 2707 compound 3	1-47
X	LEMAHIEU, RONALD A. ET AL: "Aromatic esters of 5-O-desosaminylerythronolide A oxime" JOURNAL OF MEDICINAL CHEMISTRY (1975), 18(8), 849-51 , 1975, XP002271661 page 850, compounds 3-11	1-47
X	PESTKA, SIDNEY ET AL: "Effect of erythromycin analogs on binding of carbon-14-labeled erythromycin to Escherichia coli ribosomes" ANTIMICROBIAL AGENTS AND CHEMOTHERAPY (1974), 6(4), 479-88 , 1974, XP009026503 page 481 compounds 29-31,33-43	1-47
Y	EP 0 945 459 A (TAISHO PHARMA CO LTD) 29 September 1999 (1999-09-29) the whole document	1-47
Y	EP 1 044 985 A (HOKURIKU PHARMACEUTICAL) 18 October 2000 (2000-10-18) the whole document	1-47
Y	US 6 084 079 A (PREMCHANDRAN RAMIYA H ET AL) 4 July 2000 (2000-07-04) column 1	1-47
Y	US 6 077 943 A (KATO MASAYASU ET AL) 20 June 2000 (2000-06-20) column 2 -column 6	1-47

INTERNATIONAL SEARCH REPORT

PCT/IB 03/04166

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3923784	A	02-12-1975	NONE	
EP 0945459	A	29-09-1999	AU 720921 B2	15-06-2000
			AU 4473097 A	22-06-1998
			BR 9713302 A	21-03-2000
			CA 2272620 A1	04-06-1998
			EP 0945459 A1	29-09-1999
			US 6191118 B1	20-02-2001
			CN 1238777 A	15-12-1999
			CZ 9901807 A3	15-09-1999
			HU 0000198 A2	28-08-2000
			WO 9823628 A1	04-06-1998
			ZA 9710648 A	23-06-1998
EP 1044985	A	18-10-2000	AU 1504899 A	28-06-1999
			EP 1044985 A1	18-10-2000
			CN 1284961 T	21-02-2001
			WO 9929709 A1	17-06-1999
			JP 11236395 A	31-08-1999
US 6084079	A	04-07-2000	NONE	
US 6077943	A	20-06-2000	AU 2231097 A	16-09-1997
			CA 2247929 A1	04-09-1997
			EP 0884322 A1	16-12-1998
			JP 10067795 A	10-03-1998
			WO 9731930 A1	04-09-1997